

## A STUDY ON METRICS AND STATISTICAL ANALYSIS

DĂNUȚ MARCU

**Abstract.** The purpose of this article is to introduce some classes of metrics, to describe their importance to mathematics and the sciences, to state the basic theorems concerning these classes, to state some new theorems which we have obtained by using topological methods, and even provide a proof here and there. But, the main purpose, is to state many of the open problems around these concepts and to show how much of this subject might be understood by topological means.

### 1. Introduction

Everyone is familiar with the *triangle inequality*. This inequality played a major role in the definition of a topological space.

$$\rho(a, b) \leq \rho(a, c) + \rho(b, c)$$

Still familiar to topologists is the *ultrametric inequality*.

$$\rho(a, b) \leq \max\{\rho(a, c), \rho(b, c)\}$$

But there are more inequalities of importance to mathematics which topologists are not familiar with. For example, there is the *four-point inequality*,

$$\rho(a, b) + \rho(c, d) \leq \max\{\rho(b, c) + \rho(a, d), \rho(a, c) + \rho(b, d)\}$$

and there is the *pentagon inequality*

$$\rho(a, b) + \rho(c, d) + \rho(c, e) + \rho(d, e) \leq \rho(a, c) + \rho(a, d) + \rho(a, e) + \rho(b, c) + \rho(b, d) + \rho(b, e)$$

---

Received by the editors: 25.03.2004.

2000 *Mathematics Subject Classification.* 62C07.

*Key words and phrases.* metrics, ultrametrics, statistical analysis.

and there is the *negative-type inequality*

$$\rho(a, b) + \rho(b, c) + \rho(a, c) + \rho(d, e) + \rho(d, f) + \rho(e, f)$$

$$\leq \rho(a, d) + \rho(a, e) + \rho(a, f) + \rho(b, d) + \rho(b, e) + \rho(b, f) + \rho(c, d) + \rho(c, e) + \rho(c, f)$$

All of these inequalities turn out to be important in various parts of mathematics and, especially, in the applications of mathematics to the sciences.

## 2. Statistics

The standard definition states that multivariate statistical analysis and, especially, that more applied part of multivariate statistical analysis which is called multivariate data analysis, is concerned with data collected on several dimensions of the same individual. A cursory examination of the literature of that subject reveals that a major concern, worthy of a few chapters in a typical textbook, is the following situation and resulting problem: For each of  $n$  objects, each of  $k$  tests is performed with a result which might be a real number. This gives us an  $n \times k$  matrix. We wish to combine this test data and produce an  $n \times n$  matrix of non-negative reals which measures the “similarity” or “dissimilarity” of the objects so far as their test results indicate. If the tests have been designed to give a reasonable notion of similarity, then this *similarity matrix* usually satisfies the axioms of a metric space. We wish to determine what kind of distance concept has been isolated, that is, what kind of metric space has been constructed.<sup>1</sup> Of course, with real data, things are not as simple as we have described. In most cases, the data has to be *transformed*, some data is missing and has to be *reconstructed* and the data has error or even spurious entries and has to be *approximated*. Only then can the data be *represented* in some fashion which makes it possible to use our human facilities to understand this data. So, before analysing distance data, we need some means of *classifying* metric spaces and some compendium of reasonable representations or *embeddings*.

---

<sup>1</sup>This topic is a huge one. There are many textbooks devoted to the various aspects of this problem. A bibliography listing only articles which appeared up to 1975 has 7530 entries.

### 3. Kinds of Metrics

Here is a list of the basic kinds of metrics:

- 1. ultrametric
- 2a.  $L_2$ -embeddable
- 2b. four-point property
- 3.  $L_1$ -embeddable
- 4. hypermetric
- 5. spherical
- 6. negative-type
- 7. one positive eigenvalue
- 8.  $L_\infty$ -embeddable

Each property implies those properties listed below it, except that 2a. does not imply 2b.

The purpose of this article is to introduce these classes of metrics, to describe their importance to mathematics and the sciences, to state the basic theorems concerning these classes, to state some new theorems which we have obtained by using topological methods, and even provide a proof here and there. But, the main purpose, is to state many of the open problems around these concepts and to show how much of this subject might be understood by topological means.

### 4. Ultrametries

Ultrametric spaces are well-known to topologists and perhaps even better known to number theorists and analysts. K. Hensel invented the  $p$ -adic numbers in 1897. These numbers carry a natural ultrametric structure and there are now textbooks on “Ultrametric Calculus” and “Non-Archimedean Functional Analysis”. A closely related topic which has attracted attention of many topologists is spherical completeness. The ultrametric inequality was formulated at least as early as 1934 by Hausdorff, but the term *ultrametric* was coined only in 1944 by M. Krasner. In 1956,

deGroot characterized the ultrametric spaces, up to homeomorphism, as the strongly zero-dimensional metric spaces.<sup>2</sup>

But, questions at a topological level of generality, remain open. It does not seem to be known which non-metric spaces are “essentially” ultrametric:

**Problem 1.** *Characterize those topological spaces  $X$  such that, for every continuous pseudometric  $\rho$  on  $X$ , there is a continuous ultra-pseudometric  $\sigma$  on  $X$  which generates a larger topology.*

Ultrametric spaces have emerged in the last fifteen years as a major concern in statistical mechanics, in neural networks and in optimization theory. The history of this emergence is quite interesting.

In 1984, Mezard, Parisi, Sourlas, Toulouse and Virasoro published an article on the mean-field theory of spin glasses in which they established that the distribution of “pure states” in the configuration space is an ultrametric subspace. Within a few years, it was shown that the “graph partitioning problem” in finite combinatorics could be “mapped onto” the spin glass problem and thus that the solution space for this problem also has an ultrametric structure. S. Kirkpatrick then found numerically that the solutions for certain *travelling salesman problems*<sup>3</sup> seem to scatter in an ultrametric fashion. J. P. Bouchaud and P. Le Doussal have conjectured that, in optimization problems in which “the imposed constraints cannot all be satisfied simultaneously, the optimal configurations (i.e., those which minimize the number of unsatisfied constraints) spread in an ultrametric way in the configuration space”. These kinds of problems are known as *frustrated optimization problems*.

No results of this kind have actually been proven, except in special classes of spin glasses. All other indications are numerical or by reduction. It would be of major significance to many fields to show that this phenomenon occurs under some general circumstances.

---

<sup>2</sup>Nyikos and Purisch have extensively investigated the relationship between ultrametrics and generalized metrics and orderability

<sup>3</sup>Is there an infinitary version of the travelling salesman problem? Examples might be “When do metric spaces admit space-filling curves of finite length?” or “When do they admit  $\epsilon$ -dense curves of finite length for each  $\epsilon > 0$ ?”

**Problem 2.** Give some reasonable conditions on non-negative continuous real-valued functions  $\{f_i : i < n\}$  on a metric space  $X$  so that, if  $K$  is minimal for  $Y = \{x \in X : \sum\{f_i(x) : i < n\} = K\}$  non-empty, then  $Y$  is ultrametric. Formulate this question more accurately.

A recent and effective strategy in handling optimization problems is to use simulated annealing and *random walks* to find global solutions. In problems where the local solutions have an ultrametric structure, it is therefore essential to understand random walks on ultrametric spaces. There has been much work already on different ways in which to define such random walks.

There are, undoubtedly, quite general theorems which show that the natural metric on sufficiently few independent stochastic processes which are nontrivial on sufficiently few of sufficiently many coordinates is arbitrarily close to being ultrametric. It seems likely that, to obtain a statement and proof of such a theorem, we should state and prove an infinite version first.

**Problem 3.** Let  $\{R^x : x \in X\}$  be a finite set of independent stochastic processes acting on  $\mathbb{R}^\omega$ , independently of the coordinates, so that

$$(\forall x \in X)(\forall t \in \mathbb{R}) \text{Prob}_t(|\{n \in \omega : R^x(n) \neq 0\}| < \omega) = 1$$

Let  $d$  be the metric defined on  $X$  by  $d(x, x') = E(L_1(|R^x - R^{x'}|))$  for a suitable measure on  $\omega$ . Prove that  $d$  is an ultrametric.

**Problem 4.** Can an asymptotic finitary version of problem 3 be stated and proved? Can the assumptions be made sufficiently reasonable so as to show that the numerical evidence for ultrametricity of phylogenetic trees in evolution is inevitable?

In examining the numerical evidence for ultrametricity, and in proving theoretical results about the tendency of finite data to approach ultrametricity, there is a need for answering a fundamental question: How can we measure how far a given metric is from being an ultrametric?

The main method used in spin glasses for answering this question is based on the following:

**Proposition 1 (Jardine, 1967).** *If  $\rho$  is a metric on a finite set, then there is an ultrametric  $\tau$  which minimizes  $\sup\{|\rho(x, y) - \tau(x, y)| : x, y \in X\}$  among those  $\tau$  such that  $(\forall x, y \in X)\tau(x, y) \leq \rho(x, y)$ .*

This analog of the subharmonic in potential theory which is called the subdominant ultrametric can be quite pathological. R. Rammal, G. Toulouse and M. Virasoro in their article *Ultrametricity for Physicists* ask whether there are optimal  $l_p$  ultrametric approximations for a given metric where  $1 \leq p \leq \infty$  (and specifically ask it for 1 and  $\infty$ ). Noting that the proposition can be viewed as an optimal  $l_\infty$  ultrametric approximation among those ultrametries below a given metric, we have obtained the next result:

**Theorem 1.** *If  $\rho$  is a metric on a finite space, then there is an ultrametric  $\tau$  which minimizes  $\sup\{|\rho(x, y) - \tau(x, y)| : x, y \in X\}$  among all ultrametries  $\tau$ .*

There may be several choices for the ultrametric in theorem 1 but perhaps this duplication only occurs in a trivial way.

**Problem 5.** *Is there, up to some kind of manipulation, always an unique ultrametric  $\tau$  which minimizes  $\sup\{|\rho(x, y) - \tau(x, y)| : x, y \in X\}$  among all ultrametries  $\tau$ ?*

But, our construction in theorem 1, seems to take exponential time, while Jardine's only takes polynomial time.

**Problem 6.** *Is there a polynomial algorithm for computing an ultrametric  $\tau$  which minimizes  $\sup\{|\rho(x, y) - \tau(x, y)| : x, y \in X\}$  among all ultrametries  $\tau$ ?*

Krivanek showed that computing the closest ultrametric above a given metric is NP-complete.

**Problem 7.** *Show that the subdominant ultrametric can be quite pathological. That is, show that the subdominant ultrametric of a given metric  $\rho$  can be arbitrarily close to zero, even when there is an ultrametric quite close to  $\rho$  in the supremum norm.*

Jardine's theorem was extended by Bayod and Martinez-Maurica, in 1990, to totally disconnected locally compact spaces. But, they failed to obtain a characterization.

**Problem 8.** *Characterize those metric spaces which have a subdominant ultrametric.*

**Problem 9.** *Can theorem 1 be extended to a reasonable class of infinite metric spaces?*

Returning to the problem of Rammal, Toulouse and Virasoro:

**Problem 10.** *If  $\rho$  is a metric on a finite (or arbitrary) set, then is there an ultrametric  $\tau$  which minimizes  $\sum\{|\rho(x, y) - \tau(x, y)| : x, y \in X\}$  among all ultrametries  $\tau$ ? How does one construct  $\tau$ ?*

**Problem 11.** *Which metric spaces have an (uniformly) equivalent metric  $\rho$  for which there is an ultrametric  $\tau$  such that  $\sum\{|\rho(x, y) - \tau(x, y)| : x, y \in X\}$  is finite?*

It would be quite useful to associate, to each metric, an ultrametric which is somehow derived from it in a natural way. But, this seems unlikely.

**Problem 12.** *Let the family of all metrics on a (finite, countable or arbitrary) set  $X$  be equipped with an  $l_p$  metric. Is there a continuous retraction of metrics onto ultrametries?*

Note that when  $p = \infty$ , this problem is entirely topological.

Ultrametric spaces can be embedded in linearly ordered spaces, but this is not an isometric embedding. To provide an isometric representation, we must use another device, well-known to natural scientists as a *dendrogram* (see p. 769 of Rammal). This method is equally valid for infinite spaces.

## 5. Additive Trees

The representation of ultrametries by dendrograms leads one to consider a more general kind of diagram called an additive tree in the social sciences literature or a phylogenetic tree (this term has many inexact definitions) in the biological literature. Suppose  $(V, E)$  is a tree (a graph without cycles or loops) in which each edge has an “weight” which is a non-negative real number. The distance between any two vertices  $x, y \in V$  is defined to be the sum of the weights of the edges which make up the unique minimal path from  $x$  to  $y$ . It is an exercise in graph theory to show that this distance is a metric which satisfies the four-point property.

**Theorem 2.** *Any ultrametric space satisfies the four-point property.*

In 1971, Bunemann showed that, in fact, any metric on a finite set satisfying the four-point property could be represented as the vertices of a graph equipped with this “path distance”.

**Definition 1.** *An R-tree is an (uniquely) arcwise connected metric space in which each arc is isometric to a subarc of the reals.*

In 1985, Mayer and Oversteegen constructed an universal R-tree of a given weight. This construction allows us to prove that the path metric or *intrinsic* metric on an R-tree satisfies the four-point condition and that, conversely, any metric space satisfying the four-point condition can be represented as a set of points in an R-tree.<sup>4</sup>

Indeed, the concept of an additive tree may be valuable for arbitrary completely regular spaces:

**Problem 13.** *Characterize those topological spaces  $X$  such that, for every continuous pseudometric  $\rho$  on  $X$ , there is a continuous pseudometric  $\sigma$  on  $X$  with the four-point property which generates a larger topology.*

Any linearly ordered connected compactum satisfies problem 13.

This representation by additive trees is not, by any means, only a theoretical concern. It is an useful way of representing data which satisfies the four-point condition (see p. 395 of Shepard). Note that this is the right diagram for representing evolution in which rates of evolution may be different for different species. Dendrograms assume that the rates are uniform for all species.

Additive trees are obviously easy to interpret. A topologist might ask whether one can use the intrinsic metric of more general spaces to represent metric spaces of a broader kind. The answer is yes.

**Proposition 2.** *Any separable metric space can be represented as a subset of a subspace of  $\mathbb{R}^3$  equipped with the intrinsic metric.*

---

<sup>4</sup>Rudnik and Borsuk have asked whether there is an one-dimensional subset  $X$  of  $\mathbb{R}^2$  in which every two points is joined by an arc of finite length and in which every intrinsic isometry in  $\mathbb{R}^2$  is an isometry.



But, this proposition shows by its strength, its uselessness. We must keep in mind that, to be useful, a representation must take advantage of human facilities.<sup>5</sup>

**Problem 14.** *Characterize those metric spaces which can be represented as a subset of a (simply connected) continuum in  $\mathbb{R}^2$  with the intrinsic metric.*

For example, any ultrametric space, such as  $K_5$  with the graph metric,<sup>6</sup> can be so represented but  $K(3,3)$  cannot be so represented.

**Problem 15.** *Is there a version of Kuratowski's test for planarity of graphs which answers problem 14 for graph metrics? That is, is there a finite list of "forbidden" graphs?*

While testing a metric for ultrametricity requires testing each set of three points (and thus can be done in  $O(n^3)$  computing time), testing a metric for the four-point condition seems to require testing each set of four points and that would require  $O(n^4)$  time. But, there is a beautiful way of converting additive trees into ultrametrics.

**Definition 2.** *If  $\rho$  is a metric on a set  $X$  and  $v \in X$  and  $c$  is an appropriate constant, then, for each  $x, y \in X$ , define  $\delta(x, y) = c + \rho(x, y) - \rho(x, v) - \rho(y, v)$ .  $\delta$  is the Farris transform of  $\rho$ .*

**Proposition 3 (Farris, 1970).**  *$\delta$  is an ultrametric if and only if  $\rho$  satisfies the four-point condition.*

This theorem is not hard to prove: it just requires some manipulation. Of course  $\delta$  and  $\rho$  do not generate the same topology even if we choose  $c$  carefully.

But, Farris' lemma is quite useful. We see immediately that we can test the four-point condition in just  $O(n^3)$  time. Actually, testing ultrametricity and thus the four-point condition can even be done in  $O(n^2 \log n)$  time.

**Problem 16.** *Which metric spaces can be represented up to uniform equivalence by a subset of a space (or an  $R$ -tree) with the intrinsic metric and finite total length?*

---

<sup>5</sup>But, despite this, many articles in the optimization literature ask for minimizing the total length of a graph which represents a given finite metric space. This should also be explored for infinite metric spaces.

<sup>6</sup>Any connected graph has a *graph metric* which is the largest metric in which the distance between any two vertices which are joined by an edge is 1

## 6. $L_1$ -Embeddable Metrics and their Decompositions

A metric space  $(X, \rho)$  where  $X$  is finite is said to be  $l_1$ -embeddable if we can embed  $X$  isometrically into  $l_1$ .

Do such metric spaces occur in nature? Is this class useful for statistical analysis? It is often true that real-life estimates of similarity are obtained by forming a linear combination of various criteria. Such estimates, such metrics are precisely the  $L_1$ -embeddable metrics ! Let us make this exact.

**Definition 3.** Let  $(X, \mathcal{M}, \sigma)$  be a measure space. For  $A, B \in \mathcal{M}$ , define  $\rho(A, B) = \int_{A \Delta B} d\sigma$ . We call  $\rho$   $L_1$ -embeddable.

Since we use integration, we are restricted to estimating similarity by linear combinations of various criteria. But, this still allows us to represent a broad range of metrics.

**Proposition 4.** Let  $\rho$  be a metric on a finite set. Then,  $\rho$  is  $l_1$ -embeddable if and only if  $\rho$  is  $L_1$ -embeddable.

**Theorem 3.** If a metric  $\rho$  on  $X$  satisfies the four-point-condition, then  $\rho$  is  $L_1$ -embeddable.

*Proof.* Represent  $(X, \rho)$  by a subset of an R-tree  $Y$  with the intrinsic metric. Choose  $v \in X$ . For each  $x \in X$ , let  $A_x$  be the unique shortest path in  $Y$  from  $x$  to  $v$ . Let  $\mathcal{M}$  be the set of all Borel sets of  $Y$ . Let  $\mu$  be the measure which assigns to each Borel set  $B$  the sum of the lengths of all disjoint families of paths in  $B$ . Let  $f$  be the constant one function. Now, the intrinsic metric between  $x$  and  $y$  coincides with the  $L_1$  metric on  $(Y, \mathcal{M}, \sigma)$ .

In the analysis of statistical data, it is not only important to recognize  $L_1$ -embeddable distances but also to be able to decompose distance data into an  $L_1$ -combination of more primitive distances. That is, we want to be able to carry out “linear decompositions” whenever this is possible and to identify when this is not possible.

**Definition 4.** Suppose  $(X, \rho)$  is a metric space. If there are metric spaces  $\{(X_i, \rho_i) : i \in I\}$  and a one-to-one map  $\pi : X \rightarrow \prod\{X_i : i \in I\}$  such that  $(\forall x, y \in X)\rho(x, y) = \sum_{i \in I} |\pi(x)(i) - \pi(y)(i)|$  and if  $\{\pi(x)(i) : x \in X\} = X_i$ , then we say  $\pi$  is a decomposition  $(X, \rho)$  as a subdirect  $L_1$ -product of metric spaces.

This is motivated by the important existence of subdirect representations in algebra.

**Theorem 4.** Every metric space can be decomposed in a “maximal” manner as a subdirect  $L_1$ -product of subsets of the reals and one more irreducible metric space. Every  $L_1$ -embeddable metric space is decomposed completely into a subdirect  $L_1$ -product of subsets of the reals.

*Proof.* Construct  $\pi$ , inductively, on an well-ordered set  $I$ .<sup>7</sup> If this has been done on an initial segment  $J \subset I$  and  $i$  is the least element of  $I - J$ , then define  $\rho^*(x, y) = \rho(x, y) - \sum\{|\pi(x)(i) - \pi(y)(i)| : i \in J\}$  and let  $\Sigma = \{\sigma \in \mathbb{R}^X : \rho^* - \sigma \text{ satisfies the triangle inequality}\}$  be partially ordered by defining  $\sigma \leq \sigma'$  if, for all  $x, x' \in X$ ,  $\sigma(x, x') \leq \sigma'(x, x')$ . Choose a maximal  $\sigma \in \Sigma$  and define, for each  $x \in X$ ,  $\pi(x)(i) = \sigma(i)$ .

**Problem 17.** Is there a “maximal” decomposition of metric spaces as a subdirect  $L_1$ -product of additive trees (or Hilbert spaces) and one more irreducible metric space so that every additive tree (or Hilbert space) remains its unique factor?

The notion of  $L_1$ -decomposition is well-motivated by the central importance of “dimension reduction” in multivariate data analysis. In his influential textbook, Kshirsagar said “The aim of the statistician undertaking multivariate analysis is to reduce the number of variables by employing suitable linear transformations ... thus reduces the dimensionality of the problem.” Reasonable decompositions accomplish this by removing the interaction between coordinates.

**Problem 18.** Are there reasonable  $L_p$  decompositions for  $1 < p \leq \infty$ ?

---

<sup>7</sup>The reals themselves can be decomposed into two copies of the reals, namely as the line  $y = x$ , and this is why we require an well-ordering of the factors. With a restriction to integer-valued metrics, this is no longer an issue.

A more useful  $L_1$ -decomposition would do more and break down the remaining irreducible factor in theorem 4 into an  $L_1$ -product of other irreducible factors whenever possible. We are able neither to prove such a theorem or even to formulate this accurately. The criterion by which such a decomposition should be judged is that it should have as a corollary the following result of R. L. Graham and P. M. Winkler and reported in Proc. Nat. Acad. Sci. 81 (1984) 7259.

**Theorem 5 (Graham, Winkler).** *Any finite graph can be canonically embedded isometrically into a maximum cartesian product of irreducible factors.*

The existence of the decomposition by subdirect products for varieties is a true theorem of universal algebra but, this is not a variety and so this seems to be of no help.

The general problem of identifying  $L_1$ -embeddability turns out to be significant in operations research. The problem of *multicommodity flows* is set in a graph in which each edge has a capacity and a demand. We seek a flow on the edges of the graph so that flow on each edge meets demand and does not exceed capacity. The so-called Japanese theorem of 1971 states that a capacity and demand are *feasible* i.e., can be met if there is a metric  $\rho$  on the vertices of the graph so that  $(c-r)\rho \geq 0$ . The celebrated Ford-Fulkerson theorem in operations research is just this theorem in the special and tractable case of single commodity flows in which the demand occurs on a single edge. Usually, the Ford-Fulkerson condition is not sufficient when the demand is more complicated. However, Lomonosov showed in 1978 that this condition is still sufficient when the demand lies on an  $L_1$ -embeddable subgraph.

## 7. Graphs and Hamming Distance

Indeed, theorem 5 illustrates the intimate connection between  $L_1$ -embeddability and Hamming distance. If we use factors in which all non-zero distances are 1 and a counting measure, then the  $L_1$ -distance is precisely the Hamming distance. This Hamming distance is useful in estimating distances between binary

strings, since error-correcting codes can be designed which do nothing more than replace a string with the “closest” string of a certain kind. Although Avis showed that any finite  $L_1$ -embeddable metric space embeds in a “weighted” hypercube, it is not true that an integer-valued  $L_1$ -embeddable metric can be embedded in the hypercube  $2^\kappa$  with the Hamming distance.

**Problem 19.** *Give necessary and sufficient conditions for an integer-valued ( $L_1$ -embeddable) metric to be embeddable in  $2^\kappa$  with the Hamming distance.*

For example, a necessary condition is that triangles must have even perimeter.

There is a huge literature on graphs which can be embedded in hypercubes and metrics which can be embedded in graphs<sup>8</sup>, but this beautiful theory carries us too far away from our topic.

## 8. Compactness and $L_\infty$ -Embeddable Metrics

A classical result of Banach and Mazur, published in 1932, states that any separable metric space can be isometrically embedded in  $L_\infty(\kappa)$  when  $\kappa$  is the continuum. But, more is true. Suppose  $(X, d)$  is a metric space. Fix  $a \in X$  and define an isometric embedding  $\pi$  of  $X$  into  $C^*(X) \subset L_\infty(|X|)$  by defining  $\pi(x)$  by setting  $\pi(x)(x') = d(x, x') - d(a, x')$ .

**Theorem 6 (Banach, Mazur; 1932).** *Any metric space can be isometrically embedded in  $L_\infty(\kappa)$  for sufficiently large  $\kappa$ .*

This theorem, surprisingly, is essentially finitary.

**Theorem 7.** *If every finite subset of a metric space  $X$  is  $L_\infty$ -embeddable, then  $X$  is  $L_\infty$ -embeddable.*

*Proof.* Define, for each finite  $F \subset X$ ,  $E(F)$  to be the set of all mappings  $\phi$  from  $X$  into  $\mathbb{R}^\kappa$  which are isometric when restricted to  $F$  and achieve the supremum, for any pair, on a coordinate specifically assigned to that pair. These form a centred family of closed sets. If we restrict ourselves to maps which, for some  $x \in X$ , satisfy

---

<sup>8</sup>Djoković characterized graphs that can be embedded into hypercubes in 1973.

$\phi(x) \equiv 0$ , then each  $E(F)$  is a subset of a fixed compact set and so we have a nonempty intersection.<sup>9</sup>

This leads us to the three basic compactness problems for  $L_\infty$ -embeddable (or  $L_1$ -embeddable, or  $L_p$ -embeddable) metrics.

- If every finite subset of a metric space  $X$  can be embedded in  $l_\infty$  (or  $l_1$ , or  $l_p$ ), then must  $X$  be embeddable in some  $L_\infty$  (or  $L_1$ , or  $L_p$ )?
- If  $n \in \omega$ , then what is the minimal  $k_n \in \omega$  (if it exists) such that any ( $l_1$ -embeddable,  $l_p$ -embeddable) finite metric space of size  $n$  can be embedded in  $l_\infty^{k_n}$  ( $l_1^{k_n}$ ,  $l_p^{k_n}$ )?
- If  $n \in \omega$ , then what is the minimal  $k_n \leq \omega$  (if it exists) such that any metric space which cannot be embedded in  $l_\infty^n$  ( $l_1^n$ ,  $l_p^n$ ) has a subspace of size  $k_n$  which also cannot be embedded in  $l_\infty^{k_n}$  ( $l_1^{k_n}$ ,  $l_p^{k_n}$ )?

For the first of these problems, Witsenhausen showed that, if every finite subset of a metric space  $X$  is embeddable in  $l_1$ , then  $X$  is embeddable in some  $L_1$ . Results of Yang and Zhang show that, if every finite subset of a metric space  $X$  is embeddable in  $l_2$ , then  $X$  is embeddable in some  $L_2$ . The situation for  $L_p$  seems to be unclear:

**Problem 20.** *If every finite subset of a metric space  $X$  can be embedded in  $l_p$ , then must  $X$  be embeddable in some  $L_p$ ?*

**Problem 21.** *Find a general compactness theorem which implies that the solution to the first compactness problem is positive for all  $p$ .*

For the second problem, Schoenberg noted in 1938 that, although the construction in the proof of theorem 6 above seems to require  $n$  coordinates, we can omit one coordinate without difficulty. This shows that  $k_n \leq n - 1$  for  $l_\infty$ . Wolfe showed that, in fact,  $k_n \leq n - 2$  for  $l_\infty$ . Witsenhausen has obtained the lower and upper bounds  $n - 2 \leq k_n \leq n(n - 1)/2$  for  $l_1$  and, later, Ball showed that  $k_n \leq n(n - 1)/2$  for any  $l_p$ . But, none of these results solve the problem completely:

---

<sup>9</sup>Of course,  $L_p$  might not be locally compact but this is irrelevant. We work in the Tychonoff product topology.

**Problem 22.** *If  $n \in \omega$ , then what is the minimal  $k_n \in \omega$  (if it exists) such that any  $l_1$ -embeddable finite metric space of size  $n$  can be embedded in  $l_1^{k_n}$ ? What about for  $l_p$  when  $1 < p \leq \omega$ ?*

This second problem has an interesting variation. Suppose  $D = \{1, 2, 3\}$  has the “distance” in which 1 and 3 are distance one apart and all other pairs are at distance zero. What is the least  $k_n$  such that any connected graph on  $n$  vertices can be embedded in a product of  $k_n$  many copies of  $D$  with the  $L_1$  distance? It is not obvious that  $k_n$  exists and is finite.

This may seem a strange problem, but this is exactly the “addressing problem for loop switching” posed by R. L. Graham and H. O. Pollak in 1971 in the Bell System Technical Journal and solved by P. M. Winkler in 1983. The answer is  $k_n = n - 1$ .<sup>10</sup>

The third problem is quite interesting. It may involve finite approximations to topological orientability.

**Proposition 5 (S. Malitz and J. Malitz, 1992).** *If a metric space  $X$  cannot be embedded in  $\mathbb{R}^2$  with the  $l_\infty$ -norm (or, equivalently, the  $l_1$ -norm), then  $X$  has a subspace of size 11 which cannot be embedded in  $\mathbb{R}^2$  with the  $l_\infty$ -norm (or, equivalently, the  $l_1$ -norm). Thus, determining whether a finite metric space can be embedded in  $\mathbb{R}^2$  with the  $l_\infty$ -norm can be done in polynomial time.*

They state the existence of such a number (like 11), for  $\mathbb{R}^n$  when  $n \geq 3$  is an open question, and that their methods get “wildly complicated”.

But, we have obtained the following results.

**Theorem 8.** *There is no  $N$  such that a finite metric space  $X$  cannot be embedded in  $\mathbb{R}^3$  with the  $l_\infty$ -norm if and only if  $X$  has a subspace of size  $N$  which cannot be embedded in  $\mathbb{R}^3$  with the  $l_\infty$ -norm.*

*Proof.* Use a Mobius strip in which the width of the strip is much smaller than  $N$  times the radius of the circle. Apply compactness to get a finite subset which is still sufficiently “Mobius”.

---

<sup>10</sup>These are “squashed cubes”, but the problem for graphs in ordinary cubes remains open.

**Problem 23.** *Is it true that there is no  $N$  such that a finite metric space  $X$  cannot be embedded in  $\mathbb{R}^3$  with the  $l_1$ -norm if and only if  $X$  has a subspace of size  $N$  which cannot be embedded in  $\mathbb{R}^3$  with the  $l_1$ -norm? Is this true for some  $\mathbb{R}^n$ ? Can the construction in theorem 8 be carried out in some  $\mathbb{R}^n$  with the  $l_1$ -norm?*

**Theorem 9.** *Determining whether a finite metric space  $X$  can be embedded in  $\mathbb{R}^6$  with the  $l_\infty$ -norm is NP-complete.*

*Proof.* The axes of a cube can be each be assigned one of three dimensions in exactly six ways. This assignment must be constant on the product of a cube and a line. If we join together two such products in such a way that all coordinates change, then knowing the assignment on one side of the join gives us exactly two possibilities on the other side of the join. Thus, using three more dimensions we can code the 3-colorability of graphs which is NP-complete.

**Problem 24.** *Let  $3 \leq n \leq 5$ . Is determining whether a finite metric space  $X$  can be embedded in  $\mathbb{R}^n$  with the  $l_\infty$ -norm NP-complete?*

**Problem 25.** *Is determining whether a finite metric space  $X$  can be embedded in  $\mathbb{R}^n$  with the  $l_1$ -norm NP-complete?*

## 9. $L_2$ -Embeddable Metrics

The problem of characterizing metric spaces which embed in Euclidean space of some dimension is a classical one and was solved by Menger in the 1930's. There is a book by Blumenthal entitled *Distance Geometry* and even a Mathematical Reviews section 51K devoted to this topic. But, in fact, this is an easy problem in  $\mathbb{R}^2$  with the Euclidean ( $l_2$ ) metric. For if a space embeds in  $\mathbb{R}^2$  and  $a, b, c$  are points in that space which do not satisfy the equality  $\rho(a, b) + \rho(b, c) = \rho(a, c)$  under any permutation, then  $a$  is, without loss of generality, embedded arbitrarily. Now,  $b$  is embedded on some circle centred at  $a$ , but otherwise its position is arbitrary. We deduce that  $c$  must be placed in one of two positions, but this choice is again arbitrary. But, now any further point must occupy an uniquely determined position. Thus, the position



of any point is determined uniquely once we have three points “in general position”. In the general setting of the Euclidean metric on  $\mathbb{R}^n$ , the situation is analogous.

Much of the work in distance geometry is devoted to characterizing Euclidean spaces, Banach spaces, hyperbolic spaces, inner product spaces and so forth entirely from the combinatorial properties of their metrics. But, we will not discuss here this fascinating topic and its intense activity since 1932 nor will we discuss the interesting work on the “distance-one-preserving” maps of A. D. Aleksandrov.

What is surprising and important to us is that ultrametrics are  $L_2$ -embeddable.

**Theorem 10 (Lemin, 1985; Vestfrid and Timan, 1979 for  $l_\infty$ ).** *Any ultrametric space of cardinality  $\kappa$  can be embedded isometrically in generalized Hilbert space  $\{f \in \mathbb{R}^\kappa : \sum\{f(\alpha) : \alpha \in \kappa\} < \infty\}$ .*

This requires some work.

Another surprising fact is that  $L_2$ -embeddable metrics are  $L_1$ -embeddable.

**Theorem 11.** *Any  $L_2$ -embeddable space is  $L_1$ -embeddable.*

**Problem 26.** *Give a direct proof that any  $L_2$  embeds isometrically into some  $L_1(\mu)$ . Can this be done by integration over projections onto hyperplanes of codimension 1? What happens for  $p \neq \infty$ ?*

But, the most important fact about  $L_2$ -embeddable metrics is that they are the basic notion of MDS: *multi-dimensional scaling*. This is a huge topic about which entire books have been written and for which there are many software packages being sold.

The basic purpose of MDS, the thing that these packages accomplish, is to take a set of data, either an  $n \times k$  matrix showing the results of tests or an  $n \times n$  matrix which already exhibits similarity data, and to do the best job possible in representing this data as points in the plane or in a higher-dimension Euclidean space.

There is a lot involved here. Scaling the similarity data with real numbers, reconstruction of missing and spurious data, approximation to a metric which is embeddable in some Euclidean space. The problem of reconstructing missing data is an

important one. Sippl and Scheraga, Proc. Nat. Acad. Sci. USA 83 (1986) 2283 and Schlitter 1987 in pursuit of reconstructing distance data in problems on nuclear magnetic resonance, showed that we need only a  $4 \times n$  submatrix of the distance matrix to reconstruct effectively in  $\mathbb{R}^3$  so long as the 4 points chosen are in general position.

**Problem 27.** *What happens in the reconstruction problem for the  $L_1$  or  $L_\infty$  metric?*

**Problem 28.** *If  $(X, \rho)$  is a metric space, then what are necessary and sufficient conditions on  $A \subset X^2$  so that, whenever  $\rho'$  is another metric on  $X$  such that  $\rho \upharpoonright A = \rho' \upharpoonright A$ , we must have  $\rho = \rho'$ . What if we only want  $\rho$  and  $\rho'$  to be equivalent or uniformly equivalent?*

**Problem 29.** *Find  $k(n)$  so that, if  $A$  is a metric space which can be embedded in  $l_\infty^n$ , then is there a finite set  $B \subset A$  of size  $k(n)$  such that knowing all the distances between points of  $A$  and points of  $B$  allows one to reconstruct the distance matrix.*

**Problem 30.** *Where does  $L_p$ -embeddable fit into the scheme we have given? Does ultrametric imply  $L_p$ -embeddable which implies  $L_1$ -embeddable, when  $p \neq \infty$ ? Are the classes of  $L_p$ -embeddable metrics comparable?*

## 10. Hypermetric Spaces and Spaces of Negative Type

The notion of  $L_1$ -embeddable differs greatly from additive trees and ultrametrics in that it does not seem to have a definition by a simple inequality. It is suspected that there are no simple characterizations of  $L_1$ -embeddable metrics, but this has never been established.

**Problem 31.** *Is there a first-order characterization of  $L_1$ -embeddability?*

A. Neyman showed in 1984 that there is no characterization which is a finite conjunction of inequalities. Of course, by compactness, there is an infinite conjunction of first-order formulas which characterizes  $L_1$ -embeddable.

The attempts to characterize  $L_1$ -embeddable by means of inequalities has led to some interesting inequalities which must be satisfied by any  $L_1$ -embeddable metric. These include the *hypermetric inequalities*.

**Definition 5.** A hypermetric inequality is defined for each  $b : X \rightarrow \mathbb{Z}$  such that  $\sum\{b(x) : x \in X\} = 1$  and states that  $\sum\{b(x)b(y)d(x,y) : x,y \in X\} \leq 0$ . A metric space which satisfies each hypermetric inequality is said to be a hypermetric space.

While this scheme is a little hard to understand at first, there are relatively few instances which are not satisfied automatically. In fact, the least complicated instance is accomplished by the  $b$ 's which are  $1, 1, 1, -1, -1$ . This yields the *pentagon inequality*, cited in the introduction. The easiest way to understand the hypermetric inequalities is to note that they forbid the bipartite graphs  $K(n, n + 1)$  when  $n \geq 2$ .

**Theorem 12.**  $L_1$ -embeddable metrics are hypermetric.

*Proof.* A cut pseudometric on a set  $X$  is a binary-valued pseudometric induced by any  $A \subset X$  which is defined by letting  $\rho(x, x') = 1$  iff  $|\{x, x'\} \cap A| = 1$ . Any  $L_1$ -embeddable metric is a linear combination of cut pseudometrics. Hypermetricity is clearly preserved by linear combinations. So, it suffices to show that cut pseudometrics are hypermetric. This means that we must show that, whenever  $a, b, c, d \geq 0$ , we have  $a + c - b - d = 1 \Rightarrow (a - b)(c - d) \leq 0$  which is easy.

Nevertheless, these inequalities do not characterize  $L_1$ -embeddable metrics. In 1977, Assouad and, independently, Avis in 1981, showed that the graph obtained by deleting two adjacent edges from  $K_7$  is hypermetric, but not  $L_1$ -embeddable. More sophisticated inequality schemes valid for  $L_1$ -embeddable metrics were devised by Deza and Laurent in 1992.

Despite their humble birth as approximations to  $L_1$ -embeddability, hypermetrics are significant to geometry. Consider the problem of identifying the metrics on  $\mathbb{R}^n$  which are scalar multiples of the usual metric on each straight line (these are called projective metrics). This is Hilbert's fourth problem. In 1974, Pogorolev characterized projective metrics in  $\mathbb{R}^2$ . In 1986, Szabo defined a complicated example of a projective metric on  $\mathbb{R}^3$  which does not satisfy Pogorolev's characterization. To see how hypermetrics are closely related to the fourth problem, we need a concept from convex geometry. A *zonoid* is a convex set which is arbitrarily close in the Hausdorff

metric to convex polytopes in  $\mathbb{R}^n$ . Alexander showed in 1988 that whenever the dual unit ball of a finite-dimensional normed linear space  $M$  (with a projective metric) is not zonoid, Pogorolev's characterization does not work. In 1975, Kelly proved that this problem is equivalent to determining whether the dual space of  $M$  is hypermetric. To get a projective metric on  $\mathbb{R}^3$  which does not satisfy Pogorolev's characterization, we need only a projective metric which is not hypermetric.  $L_\infty(\mathbb{R}^3)$  works!

**Problem 32.** *Does  $L_\infty(\mathbb{R}^3)$  satisfy the pentagonal inequality? Characterize the projective metrics on  $\mathbb{R}^3$  which disobey the pentagonal inequality or hypermetric inequalities (or weaker properties).*

It was proved in 1993 however by Deza, Grishukhin and Laurent, making use of Voronoi theory, that hypermetric spaces can be described by a finite list of inequalities. This is amazing, since the hypermetric scheme is infinite and does not seem to contain any redundancies. We don't know if this follows from logical considerations alone.

Another surprising aspect of the hypermetric inequalities is that, despite their failure to characterize the  $L_1$ -embeddable metrics, they do carry some power. Indeed, any hypermetric space still has some "Euclidean" structure.

Consider the example of a "distance" space consisting of the the points on the  $n$ -sphere with the metric defined by the square of the Euclidean metric. Of course, if we examine any three nearby and nearly collinear points, we see that this is not a metric space, but it certainly has many metric subspaces.

**Definition 6.** *If a metric space  $X$  can be isometrically embedded in some  $n$ -sphere with the square of the Euclidean metric, then we say that  $X$  is spherical.*

**Theorem 13 (Deza, Grishukhin, Laurent).** *Every finite hypermetric space is spherical.*

**Problem 33.** *Is any (countable, separable, arbitrary) hypermetric space isometrically embeddable in some appropriately defined  $\kappa$ -sphere? What is the correct infinitary notion of spherical?*

**Problem 34.** *There is at least an example of a spherical space which is not hypermetric?*

Note that it does not suffice to take an appropriate sphere, since this will not satisfy the triangle inequality.

Moving even further into weak properties, we can identify the *negative-type inequalities*. These are defined exactly like the hypermetric inequalities, except that we require only  $\sum\{b(x) : x \in X\} = 0$ .

**Definition 7.** *A negative-type inequality is defined for each  $b : X \rightarrow \mathbb{Z}$  such that  $\sum\{b(x) : x \in X\} = 0$  and states that  $\sum\{b(x)b(y)d(x,y) : x,y \in X\} \leq 0$ . A metric space which satisfies each negative-type inequality is said to be a space of negative-type.*

Again, it is easiest to understand the negative-type inequalities as forbidding the graph  $K(n, n)$  when  $n \geq 3$ .<sup>11</sup>

So, hypermetric spaces and spaces of negative-type are defined by analogous schemes of inequalities, and spherical spaces are characterized by embeddability in a specific Euclidean-style space. Nevertheless, spherical spaces interpolate hypermetric spaces and spaces of negative-type !

**Theorem 14 (Deza, Grishukhin).** *Every spherical space has negative-type and thus every hypermetric space has negative-type.*

Of course, metric spaces of negative-type need not be hypermetric. The graph  $K(2, 3)$  demonstrates this. This graph also answers one of the two parts of the next question, but which one?

**Problem 35.** *What is an example of a negative-type metric space which is not spherical? What is an example of a spherical space which is not hypermetric?*

In the application to Hilbert's fourth problem, we used the fact that  $L_\infty(\mathbb{R}^3)$  is not hypermetric.

**Problem 36.** *Is  $L_\infty(\mathbb{R}^3)$  of negative type? For which  $n$  is  $L_\infty(\mathbb{R}^n)$  of negative type?*

---

<sup>11</sup>One easily embeds  $K(2, 2)$  in  $\mathbb{R}^3$  and, of course,  $K_n$  is ultrametric.

The next classical result is beautiful and surprising and demonstrates immediately why spherical metrics are of negative-type.

**Theorem 15 (I. J. Schoenberg, 1938).** *A metric space is of negative-type if and only if it can be embedded in some  $\mathbb{R}^n$  with the metric which is the square of the Euclidean metric.*

Actually, in the language of linear algebra, this was first proved by Cayley !

Ponder theorem 15. It says that any metric of negative type can be squared and suddenly it is embedded in Euclidean space. But, this squaring is such a “nice” transformation ! The reason that we have not discussed the topological level of generality, since leaving additive trees becomes clear. All of these properties:  $L_2$ -embeddable,  $L_1$ -embeddable, hypermetric, spherical, negative type all coincide up to homeomorphism, up to uniform homeomorphism, even up to composition of the metric with a monotone function.

Let us call this composition a “scaling” and then be more exact.

**Definition 8.** *If  $f : [0, \infty) \rightarrow [0, \infty)$  is a function whose limit at zero is zero, then the scaling of a metric  $\rho$  by  $f$  is the function  $\rho_f$  defined by  $\rho_f(x, y) = f(\rho(x, y))$ .*

**Proposition 6.** *Any scale which is concave up preserves the triangle inequality.*

Delistathis has noted the well-known transformation  $x \rightarrow \frac{x}{1+x}$  which is used to bound metrics provides the most common example of an application of proposition 6.

The notion of scaling can be used to approach the problem of deciding how “geometric” these weaker metric concepts are.<sup>12</sup> Certainly, all separable metric spaces can be embedded by an uniform homeomorphism into Hilbert space (this was proved first by Mysiur, it seems). But, not all separable metric spaces can be embedded by a re-scaling into Hilbert space.

**Theorem 16.** *There is a separable metric space which cannot be scaled to embed in a pentagonal (and thus, Euclidean or negative-type) space.*

---

<sup>12</sup>Note that scaling preserves ultrametricity, but maybe not additive tree distances.

*Proof.* Take the bipartite graph  $K(n, n)$  for all possible choices of  $n$  and multiplied by all possible choices of positive rational numbers.

Every finite metric space has a scale which embeds it into  $l_2$  but whether one can get these scales in an uniform manner is unknown.

**Problem 37 (Maehara, 1986).** *Is there a scale which embeds all metric spaces of fixed size  $n$  (even size 5) into  $l_2$  simultaneously?*

## 11. Lipschitz Constants and Eigenvalues

Another property of a transformation weaker than uniform homeomorphism but incomparable to scaling is that of an  $\alpha$ -Lipschitz map. We say that two metrics  $\rho$  and  $\pi$  are  $\alpha$ -Lipschitz where  $\alpha \geq 1$  if every quotient  $\frac{\rho(x,y)}{\pi(x,y)}$  and its inverse is at most  $\alpha$ . Of course, two metrics are 1-Lipschitz if and only if they coincide. This notion enables us to ask whether an arbitrary metric is  $\alpha$ -Lipschitz to an Euclidean metric and so forth.

Note that the square root scaling is not  $\alpha$ -Lipschitz for any constant  $\alpha$ . So, there is no reason to expect  $L_2$ -embeddable,  $L_1$ -embeddable, hypermetric, and negative-type to be  $\alpha$ -Lipschitz for any constant  $\alpha$ .

**Proposition 7 (J. Bourgain, T. Figiel, V. Milman).** *There is a finite metric space which is not 2-Lipschitz isometrically embeddable in  $l_2$ .*

**Theorem 17.** *There is, for each  $\alpha > 2$ , a finite metric space which is not  $\alpha$ -Lipschitz to a space of negative type (or a subset of  $l_2$ ).*

Note that  $K(n, n)$  is easily shown not to be  $(\sqrt{2} - \epsilon)$ -Lipschitz isometrically embeddable in  $l_2$ .

**Problem 38.** *Is there a metric space of negative type which is not  $\alpha$ -Lipschitz isometrically embeddable to a subset of  $l_2$ ?*

In their pursuit of pathological examples in the geometry of Banach spaces, Bourgain, Milman and Wolfson did establish a Ramsey-theoretic theorem showing that in the disorder of arbitrary finite metric spaces can be found a certain amount

of “Euclidean behavior”. That is, arbitrary finite metric spaces do have fair-sized subsets which do embed into  $l_2$ .

**Theorem 18 (J. Bourgain, T. Figiel, V. Milman).** *For every  $\alpha > 1$ , there is  $C > 0$  such that every finite metric space contains a subset which is  $\alpha$ -Lipschitz embeddable in  $l_2$  and has size at least  $C \log |X|$ .*

Indeed, Bourgain, Milman and Wolfson defines their own metric inequality which says that a metric space has type 2 if there is  $\epsilon > 0$  so that, for any labelling of points by the vertices of an  $n$ -cube, the  $l_2$ -sum of the diagonals is less than  $\epsilon$  times the  $l_2$ -sum of the edges. They show that a metric space of type 2 contains copies of  $l_1^n$  up to a Lipschitz constant.

**Problem 39.** *Does type 2 fit naturally into the scheme of hypermetric and negative-type inequalities?*

**Problem 40.** *What Lipschitz constants, if any, exhibit the distinction between  $L_2$ -embeddable,  $L_1$ -embeddable, hypermetric, negative-type and one positive eigenvalue?*

Another transformation of metrics derives from the notion of a Robinsonian metric. This is a metric  $\rho$  whose underlying set admits a linear order  $\leq$  such that  $a \leq b \leq c \leq d \Rightarrow \rho(a, d) \leq \rho(b, c)$ . Thus, Robinsonian metrics are metrics which are “compatible” with a linear order. Ultrametrics are Robinsonian, but we know little more than this.

**Problem 41.** *Are additive metrics Robinsonian? Are Robinsonian metrics of negative type (or hypermetric)? What if we allow  $\leq$  to be a partial order of some kind?*

Let us now turn to eigenvalues. Suppose we are given any  $n$  points in some Euclidean space and compute the distance matrix. This matrix is symmetric and thus has all real eigenvalues. It has zero entries along the diagonal and has exactly one positive and  $n - 1$  negative eigenvalues. It turns out that if a metric has negative type, then it is still true that the distance matrix has exactly one positive eigenvalue.

**Theorem 19.** *Any metric space which is of negative type has a single positive eigenvalue.*



The existence of a single positive eigenvalue represents the weakest metric property which has so far been isolated.

**Definition 9.** *If  $(X, \rho)$  is a metric space and, for each finite  $\{a_i : i \in n\} \subset X$ , the  $n \times n$  distance matrix whose  $(i, j)$ -th entry is  $\rho(a_i, a_j)$  has exactly one positive eigenvalue, then we say that  $(X, \rho)$  has one positive eigenvalue.*

To see that this definition is reasonable, one should note that if a matrix has a particular eigenvalue, then any square submatrix also has that eigenvalue.  $K(3, 3)$  is not negative-type and, indeed, it has two positive eigenvalues.

**Problem 42.** *What are the metric spaces (of smallest cardinality) which do not have one positive eigenvalue?*

An example due to Winkler of a metric space with one positive eigenvalue which is not of negative type is the bipartite graph  $K(5, 2)$  with a single edge added between the two points on the “side” with only two points.<sup>13</sup>

**Problem 43.** *Can any metric space be scaled to have one positive eigenvalue?*

The scaling method we described (taking the square root) shows that any metric of negative type can be scaled to be Euclidean, but it is unknown what happens for metrics with one positive eigenvalue.

**Problem 44.** *Is there a metric space which has one positive eigenvalue which cannot be scaled to have negative type (equivalently, to be Euclidean)?*

**Problem 45.** *Which Tychonoff spaces have, for each continuous pseudometric, an equivalent (or generating a larger topology) continuous pseudometric with one positive eigenvalue?*

Further work has been done on investigating the characteristic polynomial of distance matrices of graphs by R. L. Graham and L. Lovász. This work is beyond the scope of this article, but, no doubt, investigating the characteristic polynomial of an arbitrary metric space would be rewarding.

---

<sup>13</sup>An elegant proof of this was given by Deza and Maehara in 1990 and Marcu in 1991.

**Problem 46.** *Is there an useful class of metric spaces strictly weaker than those with exactly one positive eigenvalue?*

## 12. Quasi-Metrics

The notion of asymmetric distances occur frequently in the literature. In optimization theory, for example, the “windy postman” problem is a version of the travelling salesman problem in which the quasi-metric represents times needed to cover a distance and so, “depending on the wind”, there is asymmetry.

Another significant application of asymmetric distances is in psychological measurement. The influential 1978 article by Cunningham explains why this is so. “There are some situations in which the direction of the dissimilarity measurement may make a difference.” He continues: “As an example, consider the case of people judging the similarity of two stimuli which differ markedly in their prominence or number of known traits”. In 1977, Tversky found that people gave a consistently higher rating when asked questions like “How similar is North Korea to Red China” than when asked questions like “How similar is Red China to North Korea”.

The notion of an additive tree and the notion of the four-point property both generalize to the asymmetric case naturally, but these generalizations do not seem to be equivalent. Bandelt in 1990 found equations which characterize the asymmetric generalization of additive trees.

Besides, these generalizations from the symmetric case, there is no available means of classifying asymmetric distances.

The distance matrices for finite subsets of a quasi-metric spaces are not symmetric and thus these matrices may have some eigenvalues which are not real.

**Problem 47.** *Do all quasi-metric spaces have an equivalent quasi-metric with all real eigenvalues?*

**Problem 48.** *Let  $X$  be a completely regular (topological) space. Is there, for every continuous quasi-metric on  $X$ , another continuous quasi-metric which generates a*

*larger topology and all of whose eigenvalues are real? What if we require these quasi-metrics to generate completely regular topologies?*

**Problem 49.** *Formulate problems whose solution would make progress towards the understanding of asymmetric distance data.*

### 13. Conclusion

The understanding of distance data is a fundamental goal of the natural and social sciences. To create this understanding, there are problems of reconstruction and approximation which are perhaps mainly problems in optimization theory and thus in linear algebra or non-linear analysis. But, the problems of transformation, representation and classification are topological problems. Although the data is finite, solving the corresponding infinitary problems gives asymptotic and efficient methods for solving the finite problems.<sup>14</sup> Moreover, finite combinatorists find all but the most graph-theoretic of these problems far too geometric or topological.<sup>15</sup> Although the use of distances suggests that this is a geometric problem, the importance of transforming the data in a non-linear manner, and the key role of approximation and reconstruction eliminates geometers from all but the most artificial and rigid of these problems. The importance of  $L_p$  in the classification may suggest that these problems lie in the territory of Banach space experts, but the absence of linearity immediately disqualifies these problems from consideration by all but the most heretical of functional analysts.

This is a problem which is directly adjacent to graph theory, optimization theory, operations research, geometry, and the theory of stochastic processes. This is a problem of immediate and great importance to communications theory, to statistical mechanics, to mathematical psychology, to mathematical taxonomy and to

---

<sup>14</sup>The importance of algorithms and complexity of computation is key to making the infinite important. If the uncountable fails, we must need enumeration and there will often be no algorithm. If the countably infinite fails, we must need to quantify over subsets and this often gives a lower bound on complexity.

<sup>15</sup>But, it seems that a large part of the theory of distances in graphs may be extended usefully, with some work, to a theory of  $L_1$ -embeddable metrics.

multivariate statistical analysis whose significance will only increase when a more sophisticated theory is developed. This is a problem whose solution can be developed by topologists.

## References

- [1] Hiroshi Maehara, Metric transforms of finite spaces and connected graphs, *Discrete Mathematics*, **61**(1986), 235–246.
- [2] Hans-Jurgen Bandelt, Recognition of tree metrics, *SIAM Journal on Discrete Mathematics*, **3**(1990), 1–6.
- [3] R. Rammal, G. Toulouse and M. A. Virasoro, Ultrametricity for physicists, *Reviews of Modern Physics*, **58**(1986), 765–788.
- [4] H. S. Witsenhausen, Minimum dimension embedding of finite metric spaces, *Journal of Combinatorial Theory, Series A*, **42**(1986), 184–199.
- [5] A. Y. Lemin, Isometric imbedding of isosceles (non-Archimedean) spaces into Euclidean ones, *Dokl. Akad. Nauk SSSR*, 285 (1985) 558–562.
- [6] H. Movahedi-Lankarani, On the inverse of Mane’s projection, *Proc. Amer. Math. Soc.*, **116**(1992), 555–560.
- [7] A. F. Timan and I. A. Vestfrid, Any separable ultrametric space is isometrically embeddable in  $l_2$ , *Funktsional Anal. i Prilozhen.*, **17**(1983), 85–86.
- [8] I. A. Vestfrid and A. F. Timan, A universality property of Hilbert spaces, *Dokl. Akad. Nauk SSSR*, **246**(1979), 528–530.
- [9] Abraham Neyman, Representation of  $L_p$ -norms and isometric embedding in  $L_p$ -spaces, *Israel J. Math.*, **48**(1984), 129–138.
- [10] John C. Mayer, Jacek Nikiel and Lex G. Oversteegen, Universal spaces for  $\mathbf{R}$ -trees, *Trans. Amer. Math. Soc.*, **334**(1992), 411–432.
- [11] J. C. Mayer, L. K. Mohler, L. G. Oversteegen and E. D. Tymchatyn, Characterization of separable metric  $\mathbf{R}$ -trees, *Proc. Amer. Math. Soc.*, **115**(1992), 257–264.
- [12] Krzysztof Rudnik, Concerning the rigidity problem for subsets of  $E^2$ , *Bull. Polish Acad. Sci. Math.*, **37**(1989), 251–254.
- [13] D. M. Avis, Hypermetric Spaces and the Hamming Cone, *Canad. J. Math.*, **33**(1981), 795–802.
- [14] Michel Deza and Monique Laurent, Facets for the cut cone. II. Clique-web inequalities, *Math. Programming, Series A*, **56**(1992), 161–188.
- [15] Michel Deza and Monique Laurent, Facets for the cut cone. I, *Math. Programming, Series A*, **56**(1992), 121–160.

- [16] M. Deza and V. P. Grishukhin, Hypermetric graphs, *Quart. J. Math. Oxford*, Series 2, **44**(1993), 399–433.
- [17] M. Deza and H. Maehara, Metric transforms and Euclidean embeddings, *Trans. Amer. Math. Soc.*, **317**(1990), 661–671.
- [18] Patrice Assouad, Sous-espaces de  $L^1$  et inegalites hypermetriques, *C. R. Acad. Sci. Paris*, Series I Math., **294**(1982), 439–442.
- [19] Patrice Assouad and Charles Delorme, Graphes plongeables dans  $L^1$ , *C. R. Acad. Sci. Paris*, Series A–B, **291**(1980), A369–A372.
- [20] Seth M. Malitz and Jerome I. Malitz, A bounded compactness theorem for  $L^1$ -embeddability of metric spaces in the plane, *Discrete Comput. Geom.*, **8**(1992), 373–385.
- [21] Ralph Alexander, Zonoid theory and Hilbert’s fourth problem, *Geom. Dedicata*, **28**(1988), 199–211.
- [22] D. Avis and V. P. Grishukhin, A bound on the  $k$ -gonality of facets of the hypermetric cone and related complexity problems, *Comput. Geom.*, **2**(1993), 241–254.
- [23] D. M. Avis, On the Extreme Rays of the Metric Cone, *Canad. J. Math.*, **32**(1980) 126–144.
- [24] R. L. Graham and P. M. Winkler, On isometric embeddings of graphs, *Trans. Amer. Math. Soc.*, **288**(1985), 527–536.
- [25] R. L. Graham and H. O. Pollak, On the Addressing Problem for Loop Switching, *Bell System Tech. J.*, **50**(1971), 2495–2519.
- [26] R. L. Graham and P. M. Winkler, Isometric embeddings of graphs, *Proc. Nat. Acad. Sci. U.S.A.*, **81**(1984), 7259–7260.
- [27] R. L. Graham and L. Lovász, Distance matrix polynomials of trees, *Problemes combinatoires et theorie des graphes*, Colloq. Internat. CNRS, Univ. Orsay, Orsay, 1976, 189–190.
- [28] R. L. Graham and L. Lovász, Distance matrix polynomials of trees, *Theory and applications of graphs*, Springer Lecture Notes in Mathematics, Vol. 642, 186–190.
- [29] Wilfried Imrich, Embedding graphs into Cartesian products, *Annals New York Acad. Sci.*, **576**(1986), 266–274.
- [30] V. D. Chepoi, Isometric subgraphs of Hamming graphs and  $d$ -convexity. *Cybernetics*, **24**(1988), 6–11.
- [31] Jolanta K. Misiewicz, Positive definite norm dependent functions on  $l^\infty$ , *Statist. Probab. Lett.*, **8**(1989), 255–260.
- [32] Marvin Marcus and Terence R. Smith, A note on the determinants and eigenvalues of distance matrices, *Linear and Multilinear Algebra*, **25**(1989), 219–230.

- [33] Pavel Krivka and Nenad Trinajstić, On the distance polynomial of a graph, *Aplikace Matematiky*, **28**(1983), 357–363.
- [34] Mirko Krivanek, The complexity of ultrametric partitions on graphs, *Inform. Process. Lett.*, **27**(1988), 265–270.
- [35] Jose M. Bayod, The space  $l^1(K)$  is not ultrametrizable, *Dekker Lecture Notes in Pure and Appl. Math.*, **137**(1992), 221–255.
- [36] Jose M. Bayod and J. Martinez-Maurica, Subdominant ultrametrics, *Proc. Amer. Math. Soc.*, **109**(1990), 829–834.
- [37] B. Van-Cutsem, Minimal superior ultrametrics under order constraint, *Discrete Math.*, **84**(1990), 201–204.
- [38] Wen-Hsiung Li, Simple method for constructing phylogenetic trees from distance matrices, *Proc. Nat. Acad. Sci. U.S.A.*, **78**(1981), 1085–1089.
- [39] Rudolf Mathar, The best Euclidian fit to a given distance matrix in prescribed dimensions, *Linear Algebra Appl.*, **67**(1985), 1–6.
- [40] Rudolf Mathar, Multidimensional scaling with constraints on the configuration, *J. Multivariate Anal.*, **33**(1990), 151–156.
- [41] W. C. Navidi, G. A. Churchill and A. von-Haeseler, Phylogenetic inference: linear invariants and maximum likelihood, *Biometrics*, **49**(1993), 543–555.
- [42] Mirko Krivanek, A note on the computational complexity of hierarchical overlapping clustering, *Aplikace Matematiky*, **30**(1985), 453–460.
- [43] David Penny, Towards a basis for classification: the incompleteness of distance measures, incompatibility analysis and phenetic classification, *Journal of Theoretical Biology*, **96**(1982), 129–142.
- [44] Lu Yang and Jing-Zhong Zhang, Metric spaces which cannot be isometrically embedded in Hilbert space, *Bull. Austral. Math. Soc.*, **30**(1984), 161–167.
- [45] Manfred J. Sippl and Harold A. Scheraga, Cayley-Menger coordinates, *Proc. Nat. Acad. Sci. U.S.A.*, **83**(1986), 2283–2287.
- [46] J. Schlitter, Calculation of coordinates from incomplete and incorrect distance data, *Z. Angew. Math. Phys.*, **38**(1987), 1–9.
- [47] Dănuț Marcu, Note on the Euclidean embeddings, *Quart. J. Math. Oxford, Series 2*, **42**(1991), 87–98.
- [48] M. Deza and P. Grishukhin, Hypermetric Graphs, *Quart. J. Math. Oxford, Series 2*, **44**(1993), 399–433.
- [49] James P. Cunningham, Free trees and bidirectional trees as representations of psychological distance, *Journal of Mathematical Psychology*, **17**(1978), 165–188.

- [50] Peter M. Winkler, Proof of the squashed cube conjecture, *Combinatorica*, **3**(1983), 135–139.
- [51] David Avis and Michel Deza, The Cut Cone,  $L^1$ -Embeddability, Complexity, and Multicommodity Flows, *Networks*, **21**(1991), 595–617.
- [52] J. P. Bouchaud and P. Le Doussal, Ultrametricity Transition in the Graph Colouring Problem, *Europhysics Letters*, **1**(1986), 91–98.
- [53] J. Douglas Carroll, Spatial, Non-Spatial and Hybrid Models for Scaling, *Psychometrika*, **41**(1976), 439–463.
- [54] David Penny, L. R. Foulds and L. M. Hendy, Testing the Theory of Evolution by Comparing Phylogenetic trees constructed from five different protein sequences, *Nature*, **297**(1982), 197–200.
- [55] A. D. Gordon, A Review of Hierarchical Classification, *J. Royal Statistical Society, Series A*, **150**(1982), 119–137.
- [56] Shmuel Sattath and Amos Tversky, Additive Similarity Trees, *Psychometrika*, **42**(1977), 319–345.
- [57] J. Bourgain, T. Figiel and V. Milman, On Hilbertian Subsets of Finite Metric Spaces, *Israel J. Math.*, **55**(1986), 147–152.
- [58] J. Bourgain, V. Milman and H. Wolfson, On Type of Metric Spaces, *Trans. Amer. Math. Soc.*, **294**(1986), 295–317.
- [59] Roger B. Shepard, Multidimensional Scaling, Tree-Fitting, and Clustering, *Science*, **210**(1980), 390–398.
- [60] John B. Kelly, Hypermetric Spaces, *Springer Lecture Notes in Mathematics*, **490**(1975), 17–31.
- [61] R. L. Graham and H. O. Pollak, On Embedding Graphs in Squashed Cubes, *Springer Lecture Notes in Mathematics*, **303**(1972), 99–110.
- [62] Nicholas Jardine and Robin Sibson, *Mathematical Taxonomy*, John Wiley and Sons, London, 1971.
- [63] Jean-Pierre Barthélemy and Alain Guénoche, *Trees and Proximity Representations*, John Wiley and Sons, Chichester, 1991.
- [64] Michael R. Anderberg, *Cluster Analysis for Applications*, Academic Press, New York, 1973.
- [65] Albert Gifi, *Nonlinear Multivariate Analysis*, John Wiley and Sons, Chichester, 1990.
- [66] E. O. Wiley, *Phylogenetics*, John Wiley and Sons, New York, 1981.
- [67] Leonard M. Blumenthal, *Theory and Applications of Distance Geometry*, Chelsea Publishing Company, Bronx, New York, 1970.

DĂNUȚ MARCU

- [68] Peter Sneath and Robert Sokal, *Numerical Taxonomy*, W. H. Freeman and Co., San Francisco, 1973.
- [69] Yadolah Dodge (editor), *Statistical Data Analysis*, North-Holland, Amsterdam, 1987.

STR. PASULUI 3, SECTOR 2, 020795-BUCHAREST, ROMANIA  
*E-mail address:* `drmarcu@yahoo.com`