

Existence and multiplicity of positive radial solutions to the Dirichlet problem for non-linear elliptic equations on annular domains

Noureddine Bouteraa and Slimane Benaicha

Abstract. In this paper, we study the existence and nonexistence of monotone positive radial solutions of elliptic boundary value problems on bounded annular domains subject to local boundary condition. By using Krasnoselskii's fixed point theorem of cone expansion-compression type we show that there exists $\lambda^* \geq \lambda_* > 0$ such that the elliptic equation has at least two, one and no radial positive solutions for $0 < \lambda \leq \lambda_*$, $\lambda_* < \lambda \leq \lambda^*$ and $\lambda > \lambda^*$ respectively. We include an example to illustrate our results.

Mathematics Subject Classification (2010): 35J25; 34B18.

Keywords: Positive solution, Elliptic equations, Existence, Multiplicity, local boundary, Green's function.

1. Introduction

In this paper, we are interested in the existence of radial positive solutions to the following, boundary value problem BVP

$$\begin{cases} -\Delta u(x) = \lambda f(|x|, u(x)), & x \in \Omega, \\ u(x) = 0, & x \in \partial\Omega, \end{cases} \quad (1.1)$$

where $\Omega = \{x \in \mathbb{R}^N : a < |x| < b\}$ with $1 < a < b$ is an annulus in \mathbb{R}^N ($N \geq 3$), $f \in C([a, b] \times [0, \infty), [0, \infty))$ and λ is a positive parameter.

The study of such problems is motivated by a lot of physical applications starting from the well-known Poisson-Boltzmann equation (see [2, 20, 30]), also they serve as models for some phenomena which arise in fluid mechanics, such as the exothermic chemical reactions or autocatalytic reactions (see [27], Section 5.11.1). The nonlinearity f in applications always has a special form and here we assume only the continuity of f and some inequalities at some points for the values of this function. However, we know that in the

integrand should stay a superposition of u with a given function (usually the exponent of u in applications) instead of u alone, but we treat this paper as the first step in this direction. The method we use is typical for local boundary value problems. We shall formulate an equivalent fixed point problem and look for its solution in the cone of nonnegative function in an appropriate Banach space. The most popular fixed point theorem in a cone is the cone-compression and cone-expansion theorem due to M. Krasnosel'skii [19] which we use in the form taken from [16]. We also point out the fact that problems of type (1) when equation does not contain paramete λ , are connected with the classical boundary value theory of Bernstein [1] (see also the studies of Granas, Gunther and Lee [15] for some extensions to nonlinear problems).

The existence and uniqueness of positive radial solutions for equations of type (1.1) when equation does not contain paramete λ , were obtained in [5], [21], [32]. Wang [32] proved that if $f : (0, \infty) \rightarrow (0, \infty)$ satisfies $\lim_{z \rightarrow 0} \frac{f(z)}{z} = \infty$ and $\lim_{z \rightarrow \infty} \frac{f(z)}{z} = 0$ then problem (1.1) when equation does not contain paramete λ , has a positive radial solution in $\Omega = \{x \in \mathbb{R}^N, N > 2\}$. That result was extended for the systems of elliptic equations by Ma [24]. We quote also the research of Ovono and Rougirel. [29] where the diffusion at each point depends on all the values of the solutions in a neighborhood of this point and Chipot et al. [12, 14]. For example in [12] considered the solvability of a class of nonlocal problems which admit a formulation in term of quasi-variational inequalities. There is a wide literature that deals with existence multiplicity results for various second-order, fourth-order and higher-order boundary value problems by different approaches, see [6, 7, 8, 9, 12, 17, 22, 23]. In 2011, Bohneure et al. [4] Studied the existence of positive increasing radial solutions for superlinear Neumann problem in the unit ball B in \mathbb{R}^N , $N \geq 2$,

$$\begin{cases} -\Delta u + u = a(|x|) f(u), & \text{in } B, \\ u > 0, & \text{in } B, \\ \partial_t u = 0, & \text{on } \partial B, \end{cases}$$

where $a \in C^1([0, 1], \mathbb{R})$, $a(0) > 0$ is nondecreasing, $f \in C^1([0, 1], \mathbb{R})$, $f(0) = 0$, $\lim_{s \rightarrow 0^+} \frac{f(s)}{s} = 0$ and $\lim_{s \rightarrow +\infty} \frac{f(s)}{s} > \frac{1}{a(0)}$.

In 2011, Hakimi and Zertiti, [17] studied the nonexistence of radial positive solutions for a nonpositone problem when the nonliearity is superlinear and has more than one zero,

$$\begin{cases} -\Delta u(x) = \lambda f(u(x)), & x \in \Omega, \\ u(x) = 0, & x \in \partial\Omega, \end{cases}$$

where $f \in C([0, +\infty), \mathbb{R})$.

In 2014, Sfecci [31], obtained the existence result by introduced the *limsup* and *liminf* types of nonresonance condition below the first positive eigenvalue for the following Neumann problems defined on the ball $B_R =$

$$\{x \in \mathbb{R}^N, |x| < R\},$$

$$\begin{cases} -\Delta u(x) = f(u(x)) + e(|x|), & \text{in } B_R, \\ u(x) = 0, & \text{on } \partial B_R, \end{cases}$$

where $f \in C(\mathbb{R}, \mathbb{R})$ and $e \in C([0, R], \mathbb{R})$.

In 2014, Butler et. al, [5] studied the positive radial solutions to the boundary value problem

$$\begin{cases} -\Delta u + u = \lambda a(|x|) f(u), & x \in \Omega, \\ \frac{\partial u}{\partial \eta} + \bar{c}(u) u = 0, & |x| = r_0, \\ u(x) \rightarrow 0, & |x| \rightarrow \infty, \end{cases}$$

where $f \in C([0, \infty), \mathbb{R})$, $\Omega = \{x \in \mathbb{R}^N : N > 2, |x| > r_0 \text{ with } r_0 > 0\}$, λ is a positive parameter, $a \in C([r_0, \infty), \mathbb{R}^+)$ such that $\lim_{r \rightarrow \infty} a(r) = 0$, $\frac{\partial}{\partial u}$ is the outward normal derivative and $\bar{c} \in C([0, \infty), (0, \infty))$.

Instead of working directly with (1.1), we note that the change of variable $u(x) = u(|x|)$, $t = |x|$ transforms (1.1) into the following boundary value problem (for details, see [14]):

$$\begin{cases} -u''(t) - \frac{N-1}{t}u'(t) = \lambda f(t, u(t)), & t \in (a, b), \\ u(a) = u(b) = 0, \end{cases}$$

where where $\lambda \geq 0$ is a positive parameter and $f \in C([a, b] \times [0, \infty), [0, \infty))$.

Inspired and motivated by the works mentioned above, we deal with existence and nonexistence of radial positive solutions to the BVP (1.1) i.e., an equivalent problem (2.1) by using of the fixed point theorem together with the properties of Green's function and we impose certain conditions on f . The paper is organized as follows. In Section 2, we present that a nontrivial and nonnegative solution of BVP (2.1) is monotone positive solution. In Section 3, we obtain some results of the existence, multiplicity and nonexistence positive solutions for BVP (2.1) depends on the parameter λ and we give an exemple to illustrate our results.

2. Preliminaries

We shall consider the Banach space $E = C[a, b]$ equipped with sup norm $\|u\| = \max_{a \leq t \leq b} |u(t)|$ and $C^+[a, b]$ is the cone of nonnegative functions in $C[a, b]$, where $1 < a < b$.

Definition 2.1. *Anonempty closed and convex set $P \subset E$ is called a cone of E if it satisfies*

- (i) $u \in P, r > 0$ implies $ru \in P$,
- (ii) $u \in P, -u \in P$ implies $u = \theta$, where θ denote the zero element of E .

Definition 2.2. *A cone P is said to be normal if there exists a positive number N called the normal constant of P , such that $\theta \leq u \leq v$ implies $\|u\| \leq N \|v\|$.*

We are interested in finding radial solutions for problem (1.1). We proceed as in introduction, setting $u(x) = u(|x|)$ $t = |x|$, we have the following equivalent boundary value problem

$$\begin{cases} -u''(t) - \frac{N-1}{t}u(t) = \lambda f(t, u(t)), & t \in (a, b), \\ u(a) = u(b) = 0. \end{cases} \quad (2.1)$$

We observe that the existence and nonexistence of radial positive solutions of (1.1) is equivalent to the existence and nonexistence of positive solutions of the problem (2.1).

In arriving our results, we need the following six preliminary lemmas. The first one is well known.

Lemma 2.3. (see, [13]) *Let $y(\cdot) \in C[a, b]$. If $u \in C^4[a, b]$, then the BVP*

$$\begin{cases} -u''(t) - \frac{N-1}{t}u(t) = y(t), & t \in (a, b), \\ u(a) = u(b) = 0, \end{cases}$$

has a unique solution

$$u(t) = \int_a^b s^{N-1} G(t, s) y(s) ds, \quad N > 2,$$

where

$$G(t, s) = \begin{cases} \frac{\left(1 - \left(\frac{a}{s}\right)^{N-2}\right) \left(\left(\frac{b}{t}\right)^{N-2} - 1\right)}{(N-2)(b^{N-2} - a^{N-2})}, & a \leq t \leq s \leq b, \\ \frac{\left(1 - \left(\frac{a}{t}\right)^{N-2}\right) \left(\left(\frac{b}{s}\right)^{N-2} - 1\right)}{(N-2)(b^{N-2} - a^{N-2})}, & a \leq s \leq t \leq b. \end{cases} \quad (2.2)$$

Lemma 2.4. *For any $(t, s) \in [a, b] \times [a, b]$, we have*

$$\frac{\left(1 - \left(\frac{a}{t}\right)^{N-2}\right)}{(N-2)(b^{N-2} - a^{N-2})} \leq G(t, s) \leq \frac{\left(\left(\frac{b}{t}\right)^{N-2} - 1\right)}{(N-2)(b^{N-2} - a^{N-2})}, \quad (2.3)$$

and

$$0 \leq \frac{\partial G}{\partial t}(t, s) \leq \frac{\left(\left(\frac{b}{s}\right)^{N-2} - 1\right) \left(\frac{(N-2)b}{a^{N-1}}\right)}{(N-2)(b^{N-2} - a^{N-2})}, \quad (t, s) \in [a, b] \times [a, b]. \quad (2.4)$$

Proof. The proof is evident, we omit it. \square

Lemma 2.5. (see [12]) *For $y(\cdot) \in C^+[a, b]$. Then the unique solution $u(t)$ of BVP*

$$\begin{cases} -u''(t) - \frac{N-1}{t}u(t) = y(t), & t \in (a, b), \\ u(a) = u(b) = 0. \end{cases}$$

is nonnegative and satisfies

$$\min_{a_1 \leq t \leq b_1} u(t) \geq c \|u\|,$$

Existence and multiplicity of positive radial solutions to the Dirichlet problem for nonlinear

where $c = \frac{\min\left\{\left(\frac{b}{b_1}\right)^{N-2} - 1, 1 - \left(\frac{a}{a_1}\right)^{N-2}\right\}}{\max\left\{\left(\frac{b}{a}\right)^{N-2} - 1, 1 - \left(\frac{a}{b}\right)^{N-2}\right\}}$ and $a_1, b_1 \in (a, b)$ with $a_1 < b_1$.

If we let

$$P = \left\{ u \in C^+[a, b] : \min_{a_1 \leq t \leq b_1} u(t) \geq c \|u\| \right\}, \quad (2.5)$$

then it is easy to see that P is a cone in $C[a, b]$. It is evident that BVP (2.1) has an integral formulation given by

$$u(t) = \lambda \int_a^b s^{N-1} G(t, s) f(s, u(s)) ds,$$

where G defined in (2.2).

Now, we define an integral operator $T_\lambda : P \rightarrow C[a, b]$ by

$$(T_\lambda u)(t) = \lambda \int_a^b s^{N-1} G(t, s) f(s, u(s)) ds.$$

Lemma 2.6. *Let $y \in C^+[a, b]$. If $u \in C^2[a, b]$ satisfies*

$$\begin{cases} u''(t) = y(t), & a \leq t \leq b, \\ u(a) = 0, \quad u(b) = 0, \end{cases}$$

then

- (i) $u(t) \geq 0$ for $t \in [a, b]$,
- (ii) $u'(t) \geq 0$ for $t \in [a, b]$.

Proof. From Lemma 2.4, we obtain $u(t) \geq 0$ and $u'(t) \geq 0$ for $t \in [a, b]$. \square

Lemma 2.7. $T_\lambda(P) \subset P$.

Proof. For any $u \in P$, we have

$$\begin{aligned} \min_{a_1 \leq t \leq b_1} T_\lambda u(t) &= \frac{\lambda}{(N-2)(b^{N-2} - a^{N-2})} \min_{a_1 \leq t \leq b_1} \left\{ \int_a^t \left(1 - \left(\frac{a}{s}\right)^{N-2}\right) s^{N-1} f(s, u(s)) \right. \\ &\times \left. \left(\left(\frac{b}{t}\right)^{N-2} - 1 \right) ds + \int_t^b \left(1 - \left(\frac{a}{t}\right)^{N-2}\right) \left(\left(\frac{b}{s}\right)^{N-2} - 1 \right) s^{N-1} f(s, u(s)) ds \right\} \\ &\geq \frac{\lambda}{(N-2)(b^{N-2} - a^{N-2})} \min_{a_1 \leq t \leq b_1} \left\{ \int_a^t \left(1 - \left(\frac{a}{s}\right)^{N-2}\right) \left(\left(\frac{b}{b_1}\right)^{N-2} - 1 \right) \right. \\ &\times \left. s^{N-1} f(s, u(s)) ds + \int_t^b \left(1 - \left(\frac{a}{a_1}\right)^{N-2}\right) \left(\left(\frac{b}{s}\right)^{N-2} - 1 \right) s^{N-1} f(s, u(s)) ds \right\} \end{aligned}$$

$$\begin{aligned}
&\geq \frac{\lambda \min \left\{ \left(\frac{b}{b_1} \right)^{N-2} - 1, 1 - \left(\frac{a}{a_1} \right)^{N-2} \right\}}{(N-2)(b^{N-2} - a^{N-2})} \min_{a_1 \leq t \leq b_1} \left\{ \int_a^t s^{N-1} f(s, u(s)) \right. \\
&\quad \times \left. \left(1 - \left(\frac{a}{s} \right)^{N-2} \right) ds + \int_t^b \left(\left(\frac{b}{s} \right)^{N-2} - 1 \right) s^{N-1} f(s, u(s)) ds \right\} \\
&= \frac{\lambda \min \left\{ \left(\frac{b}{b_1} \right)^{N-2} - 1, 1 - \left(\frac{a}{a_1} \right)^{N-2} \right\}}{(N-2)(b^{N-2} - a^{N-2})} \min_{a_1 \leq t \leq b_1} \left\{ \int_a^t \frac{\left(\frac{b}{s} \right)^{N-2} - 1}{\left(\frac{b}{s} \right)^{N-2} - 1} s^{N-1} f(s, u(s)) \right. \\
&\quad \times \left. \left(1 - \left(\frac{a}{s} \right)^{N-2} \right) ds + \int_t^b \frac{1 - \left(\frac{a}{s} \right)^{N-2}}{1 - \left(\frac{a}{s} \right)^{N-2}} \left(\left(\frac{b}{s} \right)^{N-2} - 1 \right) s^{N-1} f(s, u(s)) ds \right\} \\
&\geq \frac{\lambda \min \left\{ \left(\frac{b}{b_1} \right)^{N-2} - 1, 1 - \left(\frac{a}{a_1} \right)^{N-2} \right\}}{(N-2)(b^{N-2} - a^{N-2})} \min_{a_1 \leq t \leq b_1} \left\{ \int_a^t \frac{\left(\frac{b}{s} \right)^{N-2} - 1}{\left(\frac{b}{a} \right)^{N-2} - 1} s^{N-1} f(s, u(s)) \right. \\
&\quad \times \left. \left(1 - \left(\frac{a}{s} \right)^{N-2} \right) ds + \int_t^b \frac{1 - \left(\frac{a}{s} \right)^{N-2}}{1 - \left(\frac{a}{b} \right)^{N-2}} \left(\left(\frac{b}{s} \right)^{N-2} - 1 \right) s^{N-1} f(s, u(s)) ds \right\} \\
&\geq \frac{c\lambda}{(N-2)(b^{N-2} - a^{N-2})} \min_{a_1 \leq t \leq b_1} \left\{ \int_a^t \left(\left(\frac{b}{s} \right)^{N-2} - 1 \right) \left(1 - \left(\frac{a}{s} \right)^{N-2} \right) \right. \\
&\quad \left. s^{N-1} f(s, u(s)) ds + \int_t^b \left(1 - \left(\frac{a}{s} \right)^{N-2} \right) \left(\left(\frac{b}{s} \right)^{N-2} - 1 \right) s^{N-1} f(s, u(s)) ds \right\} \\
&= \frac{c\lambda}{(N-2)(b^{N-2} - a^{N-2})} \int_a^b \left(\left(\frac{b}{s} \right)^{N-2} - 1 \right) \left(1 - \left(\frac{a}{s} \right)^{N-2} \right) s^{N-1} f(s, u(s)) ds \\
&\geq \frac{c\lambda}{(N-2)(b^{N-2} - a^{N-2})} \max_{a \leq t \leq b} \left\{ \int_a^t \left(\left(\frac{b}{s} \right)^{N-2} - 1 \right) \left(1 - \left(\frac{a}{s} \right)^{N-2} \right) \right. \\
&\quad \times \left. s^{N-1} f(s, u(s)) ds + \int_t^b \left(1 - \left(\frac{a}{s} \right)^{N-2} \right) \left(\left(\frac{b}{s} \right)^{N-2} - 1 \right) s^{N-1} f(s, u(s)) ds \right\} \\
&\geq \frac{c\lambda}{(N-2)(b^{N-2} - a^{N-2})} \max_{a \leq t \leq b} \left\{ \int_a^t \left(\left(\frac{b}{t} \right)^{N-2} - 1 \right) \left(1 - \left(\frac{a}{s} \right)^{N-2} \right) \right. \\
&\quad \times \left. s^{N-1} f(s, u(s)) ds + \int_t^b \left(1 - \left(\frac{a}{t} \right)^{N-2} \right) \left(\left(\frac{b}{s} \right)^{N-2} - 1 \right) s^{N-1} f(s, u(s)) ds \right\}
\end{aligned}$$

Existence and multiplicity of positive radial solutions to the Dirichlet problem for nonlinear

$$= c \max_{a \leq t \leq b} T_\lambda u(t) = c \|T_\lambda u\|.$$

In other words, we find,

$$\max_{a_1 \leq t \leq b_1} T_\lambda u(t) = \|T_\lambda u\|, \quad \forall u \in P.$$

Thus, we get that $T_\lambda : P \rightarrow P$ is well defined. Moreover, it is easy to show that T_λ is completely continuous. \square

If we let

$$K = \{u \in P / u(t) \text{ is nondecreasing}\},$$

then, it is easy to show that $K \subset P$ is also a cone in E .

Lemma 2.8. $T_\lambda(P) \subset K$.

Proof. It follows from Lemma 2.6 (ii) and Lemma 2.7. \square

Lemma 2.9. $T_\lambda : K \rightarrow K$ is completely continuous.

Proof. Let $D \subset K$ is a bounded subset. Then there exists a positive constant M_1 such that

$$\|u\| \leq M_1, \quad \forall u \in D$$

Now, we shall prove that $T_\lambda(D)$ is relatively compact in K . Suppose that $(y_k)_{k \in \mathbb{N}^*} \subset A(D)$. Then there exist $(x_k)_{k \in \mathbb{N}^*} \subset D$, such that

$$y_k = Ax_k$$

Let $M_2 = \sup_{a \leq t \leq b} |f(t, u(t))|$ for all $(t, u) \in [a, b] \times [0, M_1]$. For any $k \in \mathbb{N}^*$, by Lemma 2.2, we have

$$\begin{aligned} |y_k(t)| &= |(T_n x_k)(t)| = \lambda \left| \int_a^b s^{N-1} G(t, s) f(s, x_k(s)) ds \right| \\ &\leq \lambda M_2 \int_a^b s^{N-1} G(t, s) ds \\ &\leq \frac{1}{(N-2)(b^{N-2} - a^{N-2})} \lambda M_2 \left(\left(\frac{b}{t} \right)^{N-2} - 1 \right) \int_a^b s^{N-1} ds \\ &\leq \frac{b^N - a^N}{N(N-2)(b^{N-2} - a^{N-2})} \lambda M_2 \left(\left(\frac{b}{a} \right)^{N-2} - 1 \right), \end{aligned}$$

which implies that $(y_k(t))_{k \in \mathbb{N}^*}$ is uniformly bounded.

Now, we show that A is equicontinuous. For any $u \in K$, $n \geq 2$, and $t_1, t_2 \in [a, b]$ with $|t_1 - t_2| < \delta$, we have

$$|y_k(t_1) - y_k(t_2)| = |T_\lambda u(t_1) - T_\lambda u(t_2)|$$

$$\begin{aligned} &\leq \left| \lambda \int_a^b s^{N-1} (G(t_1, s) - G(t_2, s)) f(s, x_k(s)) ds \right| \\ &\leq \lambda M_2 \int_a^b s^{N-1} |G(t_1, s) - G(t_2, s)| ds. \end{aligned}$$

It follows from the uniform continuity of Green's function G on $[a, b] \times [a, b]$, that for any $\varepsilon > 0$, we have

$$|G(t_1, s) - G(t_2, s)| \leq \frac{\varepsilon N}{\lambda(b^N - a^N)M_2}, \quad \text{for } t_1, t_2, s \in [a, b], |t_1 - t_2| < \delta.$$

Then

$$\begin{aligned} |y_k(t_1) - y_k(t_2)| &= |T_\lambda u(t_1) - T_\lambda u(t_2)| \\ &\leq \lambda M_2 \int_a^b s^{N-1} |G(t_1, s) - G(t_2, s)| ds \\ &\leq \varepsilon. \end{aligned}$$

Therefore, A is equicontinuous. By the Ascoli-Arzela Theorem, we know that A is completely continuous. \square

By Lemmas 2.8 and 2.9, we know that if $u \in P \setminus \theta$ is solution for BVP (2.1), then u is positive solution for BVP (2.1) and it is obvious from Lemma 2.8 that if $u \in P \setminus \{\theta\}$ is a solution for BVP (2.1) then $u \in K \setminus \{\theta\}$.

3. Existence and nonexistence results

In this section we will apply theorem due Krasnoselskii to study the existence, multiplicity and nonexistence of solutions for BVP (2.1) in $K \setminus \{\theta\}$.

Theorem 3.1. (See [19]) *Let E be a Banach space and $K \subset E$ be a cone in E . Assume Ω_1 and Ω_2 are open subset of E with $0 \in \Omega_1$ and $\overline{\Omega_1} \subset \Omega_2$, $T : K \cap (\overline{\Omega_2} \setminus \Omega_1) \rightarrow K$ be a completely continuous operator such that*

- (A) $\|Tu\| \leq \|u\|$, $\forall u \in K \cap \partial\Omega_1$ and $\|Tu\| \geq \|u\|$, $\forall u \in K \cap \partial\Omega_2$; or
- (B) $\|Tu\| \geq \|u\|$, $\forall u \in K \cap \partial\Omega_1$ and $\|Tu\| \leq \|u\|$, $\forall u \in K \cap \partial\Omega_2$

Then T has a fixed point in $K \cap (\overline{\Omega_2} \setminus \Omega_1)$.

We adopt the following assumptions:

(H₁) $f(t, u(t)) \in C((a, b), [0, \infty))$ is nondecreasing in $u \in [0, \infty)$ for fixed $t \in [a, b]$.

(H₂) $F_a = \int_a^b s^{N-1} f(s, 0) ds > 0$,

(H₃) $f_\infty = \lim_{u \rightarrow \infty} \min_{t \in [\frac{a}{\lambda}, b]} \frac{f(t, u)}{u} = +\infty$.

Set

$$\Lambda = \{\lambda > 0 / \text{there exists } u_\lambda \in K \setminus \{\theta\} \text{ such that } T_\lambda u_\lambda = u_\lambda\},$$

and

$$\lambda^* = \sup \Lambda.$$

Existence and multiplicity of positive radial solutions to the Dirichlet problem for nonlinear

Lemma 3.2. *Suppose that $(H_1) - (H_3)$ hold. If $\lambda' \in \Lambda$, then $(0, \lambda'] \subset \Lambda$.*

Proof. $\lambda' \in \Lambda$ means that there exists $u_{\lambda'} \in K \setminus \{\theta\}$ such that $T_{\lambda'} u_{\lambda'} = u_{\lambda'}$. Therefore, for any $\lambda \in (0, \lambda']$ we have

$$T_{\lambda} u_{\lambda'} \leq T_{\lambda'} u_{\lambda'} = u_{\lambda'},$$

Set

$$w_0 = u_{\lambda'}, w_n = T_{\lambda} w_{n-1}, n = 1, 2, \dots$$

From (H_1) , we obtain

$$w_0(t) \geq w_1(t) \geq \dots \geq w_n(t) \geq \dots \geq \frac{F_a \lambda}{(N-2)(b^{N-2} - a^{N-2})} \left(1 - \left(\frac{a}{t}\right)^{N-2}\right),$$

by Lemma 2.9 and (H_2) , $\{w_n\}$ converges to fixed point of T_{λ} in $K \setminus \{\theta\}$. Thus $(0, \lambda'] \subset \Lambda$. The proof is complete. \square

Let

$$\lambda_* < \frac{(b^{N-2} - a^{N-2})}{F_b}, \quad F_b = \int_a^b s^{N-1} f\left(s, \left(\frac{b}{a}\right)^{N-2} - 1\right) ds,$$

$$u_0(t) = \frac{\lambda F_a}{(N-2)(b^{N-2} - a^{N-2})} \left(1 - \left(\frac{a}{t}\right)^{N-2}\right), \quad v_0(t) = \left(\left(\frac{b}{t}\right)^{N-2} - 1\right),$$

and

$$F_{\infty} = \limsup_{u \rightarrow \infty} \max_{a \leq t \leq b} \frac{f(t, u)}{u}.$$

Theorem 3.3. *Suppose that $(H_1) - (H_3)$ hold. Then T_{λ} has minimal and maximal fixed point in $[u_0, v_0]$ for $\lambda \in (0, \lambda_*]$. Moreover, there exists $\lambda^* \geq \lambda_* > 0$ such that T_{λ} has at least one and has no fixed points in $K \setminus \{\theta\}$ for $0 < \lambda < \lambda^*$ and $\lambda > \lambda^*$, respectively.*

Proof. From $(H_1) - (H_3)$ and (2.3), we have $\lambda_* > 0$. For any $\lambda \in (0, \lambda_*]$, we obtain

$$\begin{aligned} (T_{\lambda} u_0)(t) &= \lambda \int_a^b s^{N-1} G(t, s) f(s, u_0(s)) ds \\ &\geq \lambda \int_a^b s^{N-1} G(t, s) f(s, u_0(a)) ds \\ &\geq \frac{\lambda}{(N-2)(b^{N-2} - a^{N-2})} \left(1 - \left(\frac{a}{t}\right)^{N-2}\right) \int_a^b s^{N-1} f(s, 0) ds \\ &\geq \frac{\lambda F_a}{(N-2)(b^{N-2} - a^{N-2})} \left(1 - \left(\frac{a}{t}\right)^{N-2}\right) = u_0(t), \end{aligned}$$

and

$$\begin{aligned}
(T_\lambda v_0)(t) &= \lambda \int_a^b s^{N-1} G(t, s) f(s, v_0(s)) ds \\
&\leq \lambda_* \int_a^b s^{N-1} G(t, s) f(s, v_0(b)) ds \\
&\leq \frac{\lambda_*}{(N-2)(b^{N-2} - a^{N-2})} \left(\left(\frac{b}{t} \right)^{N-2} - 1 \right) \int_a^b s^{N-1} f(s, v_0(b)) ds \\
&\leq \frac{\lambda_* F_b}{(N-2)(b^{N-2} - a^{N-2})} \left(\left(\frac{b}{t} \right)^{N-2} - 1 \right) = v_0(t),
\end{aligned}$$

Set

$$u_n = T_\lambda u_{n-1}, \quad v_n = T_\lambda v_{n-1}, \quad n = 1, 2, \dots,$$

then from (H_1) , we have

$$u_0(t) \leq u_1(t) \leq \dots \leq u_n(t) \leq \dots \leq v_1(t) \leq v_0(t). \quad (3.1)$$

Lemma 2.9 implies that $\{u_n\}$ and $\{v_n\}$ converge to fixed points u_λ and v_λ of T_λ , respectively.

From (3.1) it is evident that $u_\lambda, v_\lambda \in K \setminus \{\theta\}$ are the minimal fixed point and maximal fixed point of T_λ in $[u_0, v_0]$, respectively.

By the definition of λ^* , there exists a nondecreasing sequence $\{\lambda_n\}_1^{+\infty}$ such that $\lim_{n \rightarrow +\infty} \lambda_n = \lambda^*$. Let $\{u_{\lambda_n}\}_1^{+\infty}$ is bounded subset in K . Then there exists a constant $M > 0$ such that

$$\|u_{\lambda_n}\| \leq M, \quad \text{for } n \in \mathbb{N}^*,$$

which implies that $\{u_{\lambda_n}\}_1^{+\infty}$ is uniformly bounded.

Now, we show that $\{u_{\lambda_n}\}_1^{+\infty}$ is equicontinuous. For any $u_{\lambda_n} \in K$, $n \in \mathbb{N}^*$ and $t_1, t_2 \in [a, b]$, with $|t_1 - t_2| < \delta$, we have

$$\begin{aligned}
|x_{\lambda_n}(t_1) - x_{\lambda_n}(t_2)| &\leq \lambda_* \int_a^b s^{N-1} |G(t_1, s) - G(t_2, s)| f(s, M) ds \\
&\leq \lambda_* \int_a^b s^{N-1} |G(t_1, s) - G(t_2, s)| f(s, M) ds,
\end{aligned}$$

which implies that $\{x_{\lambda_n}\}_1^{+\infty}$ is equicontinuous subset in K . Consequently, by an application of the Arzela-Ascoli theorem we conclude that $\{x_{\lambda_n}\}_1^{+\infty}$ is a

Existence and multiplicity of positive radial solutions to the Dirichlet problem for nonlinear

relatively compact set in K . So, there exists a subsequence $\{x_{\lambda_{n_i}}\} \subset \{x_{\lambda_n}\}$ converging to $x^* \in K$. Note that

$$(x_{\lambda_{n_i}})(t) = \lambda_{n_i} \int_a^b s^{N-1} G(t, s) f(s, x_{\lambda_{n_i}}(s)) ds.$$

By taking the limit we have $x^*(t) = (T_{\lambda^*} x^*)(t)$. Therefore T_λ has at least one fixed point for $0 < \lambda < \lambda^*$. Finally, for T_λ has no fixed point for $\lambda > \lambda^*$. The proof is complete. \square

Theorem 3.4. *Suppose that (H_1) , (H_3) and (2.3) hold. If $(F_{+\infty} < +\infty)$, then when $F_\infty > 0$, there exists $\lambda^* \geq \frac{N(N-2)(b^{N-2}-a^{N-2})(b^N-a^N)}{F_\infty} > 0$ such that T_λ has at least one and has no fixed points in $K \setminus \{\theta\}$ for $0 < \lambda < \lambda^*$ and $\lambda > \lambda^*$, respectively. When $F_\infty = 0$, T_λ has at least one fixed points in $K \setminus \{\theta\}$ for $\lambda > 0$.*

Proof. Since $F_\infty < \infty$, for any $\varepsilon > 0$, there exists $N_0 > 0$ such that $f(t, u) \leq (F_\infty + \varepsilon)u$ for $u > N_0$, $t \in [a, b]$.

$$\text{Let } w_0(t) = N_0 \left(\left(\frac{b}{t} \right)^{N-2} - 1 \right) \text{ and } \lambda_0 = \frac{N(N-2)(b^{N-2}-a^{N-2})(b^N-a^N)}{\left(\left(\frac{b}{a} \right)^{N-2} \right) (F_\infty + \varepsilon)},$$

then $\lambda_0 > 0$ and

$$\begin{aligned} (T_{\lambda_0} w_0)(t) &= \lambda_0 \int_a^b s^{N-1} G(t, s) f(s, w_0(s)) ds \\ &\leq \frac{\lambda_0}{(N-2)(b^{N-2}-a^{N-2})} \left(\left(\frac{b}{t} \right)^{N-2} - 1 \right) \int_a^b s^{N-1} (F_\infty + \varepsilon) w_0(s) ds \\ &\leq \frac{\lambda_0 w_0(t) (F_\infty + \varepsilon)}{(N-2)(b^{N-2}-a^{N-2})} \left(\left(\frac{b}{t} \right)^{N-3} - 1 \right) \int_a^b s^{N-1} ds \\ &\leq \frac{\lambda_0 w_0(t) (F_\infty + \varepsilon)}{N(N-2)(b^{N-2}-a^{N-2})(b^N-a^N)} \left(\left(\frac{b}{a} \right)^{N-2} - 1 \right) \\ &\leq w_0(t), \end{aligned}$$

Now, set $w_0(t) = N_0 \left(\left(\frac{b}{t} \right)^{N-2} - 1 \right)$,

$$w_n = T_{\lambda_{n-1}} w_{n-1}, \quad n = 1, 2, \dots$$

From (H_1) , we obtain

$$w_0(t) \geq w_1(t) \geq \dots \geq w_n(t) \geq \dots \geq \frac{F_a \lambda}{(N-2)(b^{N-2}-a^{N-2})} \left(1 - \left(\frac{a}{t} \right)^{N-2} \right). \quad (3.2)$$

Therefore, the sequence $\{w_n\}$ is bounded in $K \setminus \{\theta\}$. By Lemma 2.9 and the definition of λ^* , the operator T_{λ_n} completely continuous. Hence the sequence

$\{w_n\}$ is compact in $K \setminus \{\theta\}$, its also monotone. Then it is uniformly convergent to fixed points u^* of T_{λ_n} in $K \setminus \{\theta\}$. When we pass to the limit we get

$$u^* = T_{\lambda^*} u^*$$

For $\lambda > \lambda^*$, there exists $\{\lambda_n\}_1^\infty$, with $\lim_{n \rightarrow \infty} \lambda_n = \lambda$, we prove that problem has no positive solution. suppose the contrary that the problem has a positive solution x_{λ_n} , then we get

$$\begin{aligned} \|u_{\lambda_n}\| &= (T_{\lambda_n} u_{\lambda_n}) \left(\left(\frac{b}{a} \right)^{N-2} - 1 \right) \\ &\leq \frac{\lambda_n}{(N-2)(b^{N-2} - a^{N-2})} \left(\left(\frac{b}{a} \right)^{N-2} - 1 \right) \int_a^b s^{N-1} f(s, u_{\lambda_n}(s)) ds \\ &\leq \frac{\lambda_n}{(N-2)(b^{N-2} - a^{N-2})} \left(\left(\frac{b}{a} \right)^{N-2} - 1 \right) \int_a^b s^{N-1} (F_\infty + \epsilon) u_{\lambda_n}(b) ds \\ &\leq \frac{\lambda_n (b^N - a^N)}{N(N-2)(b^{N-2} - a^{N-2})} \left(\left(\frac{b}{a} \right)^{N-2} - 1 \right) (F_\infty + \epsilon) u_{\lambda_n}(b) \\ &\leq \frac{\lambda_n (b^N - a^N)}{N(N-2)(b^{N-2} - a^{N-2})} \left(\left(\frac{b}{a} \right)^{N-2} - 1 \right) (F_\infty + \epsilon) \|u_{\lambda_n}\| < \|u_{\lambda^*}\|. \end{aligned}$$

Taking the limit we obtain

$$\|u_\lambda\| < \|u_\lambda\|,$$

which is a contradiction. The proof is complete. \square

Lemma 3.5. Assume that (H_1) , (H_2) and (H_3) hold. If Λ is nonempty, then
(i) Λ is bounded from above, that $\lambda^* < +\infty$.
(ii) $\lambda^* \in \Lambda$.

Proof. Suppose to the contrary that there exists an increasing sequence $\{\lambda_n\}_1^{+\infty} \subset \Lambda$ such that $\lim_{n \rightarrow +\infty} \lambda_n = +\infty$. Set $x_{\lambda_n} \in K \setminus \{\theta\}$ is a fixed point of T_{λ_n} that is ,

$$T_{\lambda_n} u_{\lambda_n} = u_{\lambda_n}.$$

There are two cases to be considered.

Case 1. $\{u_{\lambda_n}\}_1^{+\infty}$ is bounded, that is there exists a constant $M > 0$ such that

$$\|u_{\lambda_n}\| \leq M, \text{ for } n = 1, 2, \dots$$

Hence, from (H_1) , (H_2) , and (H_3) and Lemma 2.3, we have

$$\begin{aligned} M &\geq \|u_{\lambda_n}\| \geq (T_{\lambda_n} u_{\lambda_n})(t) \\ &\geq \frac{\lambda_n}{(N-2)(b^{N-2} - a^{N-2})} \left(1 - \left(\frac{a}{b} \right)^{N-2} \right) \int_a^b s^{N-1} f(s, 0) ds \end{aligned}$$

Existence and multiplicity of positive radial solutions to the Dirichlet problem for nonlinear

$$= \frac{\lambda_n}{(N-2)(b^{N-2} - a^{N-2})} \left(1 - \left(\frac{a}{b}\right)^{N-2}\right) F_a \rightarrow +\infty,$$

which is a contradiction.

Case 2. $\{u_{\lambda_n}\}_1^{+\infty}$ is unbounded, that is there exists subsequence of $\{u_{\lambda_n}\}_1^{+\infty}$ still denoted by $\{u_{\lambda_n}\}_1^{+\infty}$ such that $\lim_{n \rightarrow +\infty} \|u_{\lambda_n}\| = +\infty$.

When (H_3) , take $L > \frac{N(N-2)(b^{N-2} - a^{N-2})}{(1 - (\frac{a}{b})^{N-2})\lambda_1}$ there exists $N_1 > 0$ such that

$f(t, u) \geq Lu$, for $u \geq N_1$, $t \in [a, b]$. Choose n_1 such that $\|u_{\lambda_{n_1}}\| > NN_1$.

Thus, for $t \in [a, b]$, we have

$$f\left(t, \frac{1}{N} \|u_{\lambda_{n_1}}\|\right) \geq \frac{1}{N} L \|u_{\lambda_{n_1}}\|.$$

Moreover, from (H_1) and the definition of K , we have

$$\begin{aligned} \|x_{\lambda_{n_1}}\| &\geq (T_{\lambda_{n_1}} u_{\lambda_{n_1}})(t) \\ &\geq \frac{\lambda_{n_1}}{(N-2)(b^{N-2} - a^{N-2})} \left(1 - \left(\frac{a}{b}\right)^{N-2}\right) \int_a^b s^{N-1} f(s, u_{\lambda_{n_1}}(s)) ds \\ &\geq \frac{\lambda_{n_1}}{(N-2)(b^{N-2} - a^{N-2})} \left(1 - \left(\frac{a}{b}\right)^{N-2}\right) \int_a^b s^{N-1} f\left(s, \frac{1}{6} \|u_{\lambda_{n_1}}\|\right) ds \\ &= \frac{\lambda_{n_1} L \left(1 - \left(\frac{a}{b}\right)^{N-2}\right)}{N(N-2)(b^{N-2} - a^{N-2})} \|u_{\lambda_{n_1}}\| > \|u_{\lambda_{n_1}}\|, \end{aligned}$$

which is a contradiction.

Consequently, we find that Λ is bounded from above.

(ii) From the definition of λ^* , there exists a nondecreasing sequence $\{\lambda_n\}_1^{+\infty}$ such that $\lim_{n \rightarrow +\infty} \lambda_n = \lambda^*$. Let $\{u_{\lambda_n}\}_1^{+\infty} \in K \setminus \{\theta\}$ be a fixed point of T_{λ_n} . Arguing similarly as above in Case 2, we can show that $\{u_{\lambda_n}\}_1^{+\infty}$ is bounded subset in K , that is there exists a constant $M > 0$. Hence from (H_1) , (H_2) , and (H_3) , we have

$$\begin{aligned} \|u_{\lambda_n}\| &= (T_{\lambda_n} u_{\lambda_n}) \left(\left(\frac{b}{a}\right)^{N-2} - 1 \right) \\ &\leq \frac{\lambda_n}{(N-2)(b^{N-2} - a^{N-2})} \left(\left(\frac{b}{a}\right)^{N-2} - 1 \right) \int_a^b s^{N-1} f(s, u_{\lambda_n}(s)) ds \\ &\leq \frac{\lambda_n}{(N-2)(b^{N-2} - a^{N-2})} \left(\left(\frac{b}{a}\right)^{N-2} - 1 \right) \int_a^b s^{N-1} f(s, u_{\lambda_n}(b)) ds \\ &\leq \frac{\lambda_n}{(N-2)(b^{N-2} - a^{N-2})} \left(\left(\frac{b}{a}\right)^{N-2} - 1 \right) \int_a^b s^{N-1} f(s, v_{\lambda_n}(b)) ds \end{aligned}$$

$$\begin{aligned}
&= \frac{\lambda_n}{(N-2)(b^{N-2} - a^{N-2})} \left(\left(\frac{b}{a} \right)^{N-2} - 1 \right) \int_a^b s^{N-1} f(s, 0) ds \\
&= \frac{\lambda_n F_a}{(N-2)(b^{N-2} - a^{N-2})} \left(\left(\frac{b}{a} \right)^{N-2} - 1 \right) \rightarrow \frac{\lambda_* F_a \left(\left(\frac{b}{a} \right)^{N-2} \right)}{(N-2)(b^{N-2} - a^{N-2})} = M,
\end{aligned}$$

as $n \rightarrow \infty$.

Therefore

$$\|u_{\lambda_n}\| \leq M, \quad n = 1, 2, \dots$$

which shows that $\{u_{\lambda_n}\}_1^{+\infty}$ is uniformly bounded.

From the proof of Theorem 3.3 we know that $\{u_{\lambda_n}\}_1^{+\infty}$ is equicontinuous subset in K and by an application of the Arzela-Ascoli theorem we conclude that $\{u_{\lambda_n}\}_1^{+\infty}$ is a relatively compact set in K . So, there exists a subsequence $\{u_{\lambda_{n_i}}\} \subset \{u_{\lambda_n}\}$ converging to $u^* \in K$. Note that

$$(u_{\lambda_{n_i}})(t) = \lambda_{n_i} \int_0^1 s^{N-1} G(t, s) f(s, u_{\lambda_{n_i}}(s)) ds.$$

By taking the limit we have

$$u^*(t) = (T_{\lambda^*} u^*)(t) \geq \frac{\lambda_1 F_a}{(N-2)(b^{N-2} - a^{N-2})} \left(1 - \left(\frac{a}{t} \right)^{N-2} \right),$$

that is $\lambda^* \in \Lambda$. The proof is complete. \square

Theorem 3.6. *Suppose that $(H_1) - (H_3)$ holds. Then there exists $\lambda^* \geq \lambda_* > 0$ such that BVP (2.1) has at least two, one and no positive solutions for $0 < \lambda \leq \lambda_*$, $\lambda_* < \lambda \leq \lambda^*$ and $\lambda > \lambda^*$ respectively.*

Proof. From (H_1) , (H_2) and (H_3) we have $(0, \lambda_*) \subset \Lambda$. So $\lambda^* \geq \lambda_* > 0$.

From Lemma 3.2 and 3.5, we have $(0, \lambda^*] = \Lambda$. Therefore, from the definition of λ^* we only to prove that T_λ has at least two fixed points in $K \setminus \{\theta\}$ for $\lambda \in (0, \lambda_*]$.

Now, given $\lambda \in (0, \lambda_*]$. Theorem 3.3 means that T_λ has at least one fixed point $u_{\lambda,1} \in K \setminus \{\theta\}$ which satisfies $\|u_{\lambda,1}\| \leq \left(\frac{b}{a} \right)^{N-2} - 1$.

Let

$$K_1 = \left\{ x \in K \mid \|u\| < \left(\frac{b}{a} \right)^{N-2} - 1 \right\}.$$

For $t \in [a, b]$, so for $u \in K$ with $\|u\| = \left(\frac{b}{a} \right)^{N-2} - 1$, i.e $u \in \partial K_1$, we have

$$\begin{aligned}
\|u\| &= \|T_\lambda u\| = (T_\lambda u) \left(\left(\frac{b}{a} \right)^{N-2} - 1 \right) \\
&\leq \frac{\lambda}{(N-2)(b^{N-2} - a^{N-2})} \left(\left(\frac{b}{a} \right)^{N-2} - 1 \right) \int_a^b s^{N-1} f(s, u(s)) ds
\end{aligned}$$

Existence and multiplicity of positive radial solutions to the Dirichlet problem for nonlinear

$$\begin{aligned}
 &\leq \frac{\lambda_*}{(N-2)(b^{N-2}-a^{N-2})} \left(\left(\frac{b}{a} \right)^{N-2} - 1 \right) \int_a^b s^{N-1} f \left(s, \left(\frac{b}{a} \right)^{N-2} - 1 \right) ds \\
 &< \frac{\left(\left(\frac{b}{a} \right)^{N-2} - 1 \right)}{N-2} < \|u\|.
 \end{aligned} \tag{3.3}$$

When (H_3) , take $L > \frac{N(N-2)(b^{N-2}-a^{N-2})}{\left(1-\left(\frac{a}{b}\right)^{N-2}\right)\lambda_1}$ there exists $N_1 > 0$ such that $f(t, u) \geq Lu$, for $u \geq N_1$, $t \in [a, b]$.

Set $K_2 = \{u : \|u\| < NN_1\}$. Then $\overline{K_1} \subset K_2$. If $u \in \partial K_2$, we have

$$\begin{aligned}
 \|u\| &= \|T_\lambda u\| = (T_\lambda u) \left(\left(\frac{b}{a} \right)^{N-2} - 1 \right) \\
 &\geq \frac{\lambda}{(N-2)(b^{N-2}-a^{N-2})} \left(1 - \left(\frac{a}{b} \right)^{N-2} \right) \int_a^b s^{N-1} f(s, u(s)) ds \\
 &\geq \frac{\lambda}{(N-2)(b^{N-2}-a^{N-2})} \left(1 - \left(\frac{a}{b} \right)^{N-2} \right) \int_a^b s^{N-1} f \left(s, \frac{1}{N} \|u\| \right) ds \\
 &\geq \frac{\lambda}{(N-2)(b^{N-2}-a^{N-2})} \left(1 - \left(\frac{a}{b} \right)^{N-2} \right) \int_a^b s^{N-1} f \left(s, \frac{1}{N} \|u(s)\| \right) ds \\
 &\geq \frac{\lambda L \left(1 - \left(\frac{a}{b} \right)^{N-2} \right)}{N(N-2)(b^{N-2}-a^{N-2})} \|u\| > \|u\|.
 \end{aligned}$$

Consequently, Applying Theorem 3.1 that T_λ has a fixed point $u_{\lambda,2} \in \overline{K_2} \setminus K_1$. Equation (3.3) implies that T_λ has no fixed point in ∂K_1 . In conclusion, for $\lambda \in (0, \lambda_*]$, T_λ has at least two fixed points $u_{\lambda,1}$ and $u_{\lambda,2}$ in K . The proof is complete. \square

We present an example to illustrate the applicability of the results shown before.

Example 3.7. Consider in \mathbb{R}^3 the elliptic boundary value problem

$$\begin{cases} -\Delta u(x) = \lambda(|x| + u + \ln(1 + u)), & x \in \Omega, \\ u(x) = 0, & x \in \partial\Omega, \end{cases} \tag{3.4}$$

To the system (3.4) we associate the the second order boundary value problem

$$\begin{cases} -u''(t) = \lambda(t + u + \ln(1 + u)), & t \in [a, b], \\ u(a) = u(b) = 0, \end{cases}$$

By direct computation, we have

$$F_\infty = 2, F_0 = \frac{1}{4}, F_1 = \frac{1}{2} + \frac{2}{3}(1 + \ln(2)) \text{ and } \lambda_* = \frac{48 - 9\pi}{6 + 8(1 + \ln(2))}.$$

So, the assumptions (H_1) , (H_2) and (H_3) are satisfied, it follows from Theorem 3.4 there exists $\lambda^* = 3 \geq \lambda_*$ such that boundary value problem (3.4) has at least one positive solution for $0 < \lambda \leq 3$ and has no positive solution for $\lambda > \lambda^*$.

References

- [1] Bernstein. S., Sur les equations du calcul des variations, Ann. Sci. Ecole Norm. Sup, 29 (1912), 431-485.
- [2] Bebernes. J, Lacey. A. A., Global existence and nite-time blow-up for a class of nonlocal parabolic problems, Adv. Dierential Equations, 2(6) (1997), 927-953.
- [3] Bandle. C, Coffman. C. V and Marcus. M., Nonlinear elliptic problems in the annulus, J. Differential Equations, 69 (1978), 322-345.
- [4] Bonheure. D and Serra. E., Multiple positive radial solutions on annulu for nonlinear Neumann problems with large growth, Nonlinear Differ. Equ. Appl, 18 (2011), 217-235.
- [5] Butler. D, Ko. E, Kyuong. E and Shivaaji. R., Positive radial solutions for elliptic equations on exterior domains with nonlinear boundary conditions, Communication on Pure and Applied Analysis, Volume 13, Number 6, (2014), 2713-27631.
- [6] Bouteraa. N and Benaicha. S., Triple positive solutions of higher-order nonlinear boundary value problems, Journal of Computer Science and Computational Mathematics, Volume 7, Issue 2, June 2017.
- [7] Bouteraa. N and Benaicha. S, Djourdem. H and Elarbi Benatia. M., Positive solutions for fourth-order two-point boundary value problem with a parameter, Romanian Journal of Mathematic and Computer Science, 2018, Volume 8, Issue 1 (2018), p. 17-30.
- [8] Bouteraa. N and Benaicha. S., Nonlinear boundary value problems for higher-order ordinary differential equation at resonance, Romanian Journal of Mathematic and Computer Science, Volume 8, Issue 2 (2018), p. 83-91.
- [9] Bouteraa. N and Benaicha. S., Existence of solutions for third-order three-point boundary value problem, Mathematica, 60 (83), No 1, 2018, pp. 12-22.
- [10] Chipot. M and Rodriguez. J. F., On a class of nonlocal nonlinear elliptic problems, Math. Modl, Nume. Anal, 26, 3 (1992), 447-468.
- [11] Chipot. M and Lovat. B., Existence and uniqueness results for a class of non-local elliptic and parabolic problems, Dyn. Contin. Discrete Impuls. Syst. Ser. A Math. Anal. 8 (2001) 35-51.
- [12] Cianciaruso. F, Infante. G and Pietramala., Solutions of perturbed Hammerstein integral equations with applications, Nonl. Anal. Real World Appl, 33 (2017), 317-347.
- [13] David. F., Radial solutions of an elliptic system, Houston J. Math, 15(1989), pp. 425-458.
- [14] Egorov. Y and Kondratiev. V., On Spectral Theory of Elliptic Operators, Birkhauser, Basel, Boston, Berlin, 1996.
- [15] Granas. A, Gunther. R and Lee. J., On a theorem of S. Bernstein, Pacic J. Math. 74 (1978), 67-82.

- [16] Guo. D and Lakshmikantham. V., *Nonlinear Problems in Abstract Cones*, Acad. Press, Inc., Boston, MA, 1988.
- [17] Hakimi. S and Zertiti. A., Nonexistence of radial positive solutions for a non-positone problem, *Elec. J. Diff. Equ*, 26 (2011), 1-7.
- [18] Infante. G and Pietramala. P., Nonzero radial solutions for a class of elliptic systems with nonlocal BCs on annular domains, *NoDEA Nonlinear Dierential Equations Appl*, 22 (2015), 979-1003.
- [19] Krasnoselskii, M., *Positive Solutions of Operator Equations*, P. Noordho Ltd., Groningen, 1964.
- [20] Krzywicki. A and Nadzieja. T., Nonlocal elliptic problems. Evolution equations: existence, regularity and singularities, *Banach Center Publ*, 52 (2000), Polish Acad. Sci., Warsaw, 147-152.
- [21] Lions. P. L., On the existence of positive solutions of semilinear elliptic equations, *SIAM Rev*, 24 (1982), 441-467.
- [22] McLeod. K and Serrin. J., Uniqueness of positive radial solutions of $\Delta u + f(u) = 0$ in \mathbb{R}^n , *Arch. Rational Mech. Anal*, 99(1987), pp. 115-145.
- [23] Marcus. M and Bandle. C., The positive radial solutions of a class of semilinear elliptic equations, *J. Reine Angrew. Math*, 401(1989), 25-59.
- [24] Ma. R., Existence of positive radial solutions for elliptic systems, *J. Math. Anal. Appl*, 201 (1996), 375-386.
- [25] Marcos do O. J, Lorca. S, Sanchez. J and Ubilla. P., Positive solutions for some nonlocal and nonvariational elliptic systems, *Comp. Variab. Ellip. Equ*, (2015), 18 pages.
- [26] Ni. W. M, Nussbaum. R. D., Uniqueness and nonuniqueness for positive radial solutions of $\Delta u + f(u, r) = 0$. *Commun. Pure Appl. Math*, 38(1), 67-108 (1985)
- [27] Ockedon. J, Howison. S, A. Lacey and Movchan. A., *Applied Partial Differential Equations*, Oxford University Press, 2003.
- [28] Ovono. A and Rougirel. A., Elliptic equations with diffusion parameterized by the range of nnonlocal interactions, *Annali di Matematica*, 009-0104-y (2009).
- [29] Precup. R, Existence., localization and multiplicity results for positive radial solutions of semilinear elliptic systems, *J. Math. Anal. Appl*, 352(2009), 48-56.
- [30] Stanczy. R., Nonlocal elliptic equations, *Nonlinear Anal*, 47 (2001), 3579-3584.

Noureddine Bouteraa

“Laboratory of Fundamental and Applied Mathematics of Oran (LMFAO)”

University of Oran1, Ahmed Benbella,

Algeria

e-mail: bouteraa-27@hotmail.fr

Slimane Benaicha

“Laboratory of Fundamental and Applied Mathematics of Oran (LMFAO)”

University of Oran1, Ahmed Benbella,

Algeria

e-mail: slimanebenaicha@yahoo.fr