

POSITIVE SOLUTIONS FOR FRACTIONAL DIFFERENTIAL EQUATIONS WITH NON-SEPARATED TYPE NONLOCAL MULTI-POINT AND MULTI-TERM INTEGRAL BOUNDARY CONDITIONS

H. DJOURDEM, S. BENAICHA

ABSTRACT. In this paper, we investigate a class of nonlinear fractional differential equations that contain both the multi-term fractional integral boundary condition and the multi-point boundary condition. By the Krasnoselskii fixed point theorem we obtain the existence of at least one positive solution. Then we obtain the existence of at least three positive solutions by the Legget-Williams fixed point. Examples are given to illustrate our main results.

1. INTRODUCTION

Differential equations of fractional order is one of the fast growing area of research in the field of mathematics and have recently been proved to be valuable tools in the modeling of many phenomena in various fields of science and engineering. Indeed, one can find numerous applications of fractional order differential equations in viscoelasticity, electro-chemistry, control theory, movement through porous media, electromagnetics, and signal processing of wireless communication system, etc (see [6, 7, 9, 10, 16, 20, 21, 23, 27, 28]). Now, there are many papers dealing with the problem for different kinds of boundary value conditions such as multi-point boundary condition (see [1, 12, 13, 14, 19, 22, 25, 29]), integral boundary condition (see [2, 4, 5, 8, 15, 30, 31]), and many other boundary conditions (see [3, 11, 18, 29, 33]).

In this paper, we are dedicated to considering fractional differential equations that contain both the multi-term fractional integral boundary condition and the multi-point boundary condition:

$$\begin{cases} D^q u(t) + f(t, u(t)) = 0, & 1 < q \leq 2, 0 < t < 1, \\ u(0) = 0, \quad u(1) = \sum_{i=1}^m \alpha_i (I^{p_i} u)(\eta) + \sum_{i=1}^m \beta_i u(\xi_i), \end{cases} \quad (1.1)$$

where D^q is the standard Riemann-Liouville fractional derivative of order q , I^{p_i} is the Riemann-Liouville fractional integral of order $p_i > 0$, $i = 1, 2, \dots, m$, $0 < \xi_1 < \xi_2 < \dots < \xi_m < 1$, $f : [0, 1] \times [0, \infty) \rightarrow [0, \infty)$ and $\alpha_i, \beta_i \geq 0$ with $i = 1, 2, \dots, m$, are real constants such that $\Gamma(q) \sum_{i=1}^m \frac{\alpha_i \eta^{p_i+q-1}}{\Gamma(p_i+q)} + \sum_{i=1}^m \beta_i \xi_i^{q-1} < 1$.

Zhou and Jiang [34] considered the fractional boundary value problem

$$\begin{cases} D_{0+}^\alpha u(t) + f(t, u(t)) = 0, & 0 < t < 1, \\ u'(0) - \beta u(\xi) = 0, \quad u'(1) + \sum_{i=1}^{m-3} \gamma_i u(\eta_i) = 0 \end{cases}$$

2010 *Mathematics Subject Classification.* 34A08, 34B15, 34B18.

Key words and phrases. fractional differential equations; Riemann-Liouville fractional derivative; multi-term fractional integral boundary condition; fixed point theorems

where α is a real number with $1 < \alpha \leq 2$, $0 \leq \beta \leq 1$, $0 \leq \gamma_i \leq 1$, $i = 1, 2, \dots, m - 3$, $0 \leq \xi < \eta_1 < \eta_2 < \dots < \eta_{m-3} \leq 1$, D_{0+}^α is the Caputo's derivative. The authors used the fixed point index theory and Krein-Rutman theorem to obtain the existence results.

Ji et al. [26] investigated the existence and multiplicity results of positive solutions for the following boundary value problem:

$$\begin{cases} D_{0+}^\alpha u(t) + f(t, u(t), D_{0+}^\mu u(t)) = 0, & 0 < t < 1, \\ u(0) = 0, \quad u(1) + D_{0+}^\beta u(1) = ku(\xi) + lD_{0+}^\beta u(\eta), \end{cases}$$

where D_{0+}^α is the Riemann-Liouville fractional derivative of order $1 < \alpha \leq 2$, $0 \leq \beta \leq 1$, $\xi, \eta \in (0, 1)$, $0 \leq \mu < 1$, $1 \leq \alpha - \beta$, $1 \leq \alpha - \mu$, $1 - l\eta^{\alpha-\beta-1}$, and $f : [0, 1] \times [0, +\infty) \times (-\infty, +\infty) \rightarrow [0, +\infty)$ is continuous. They used the Leggett-Williams fixed point theorem to obtain the existence and multiplicity results of positive solutions.

Wang et al. [32] considered the following boundary value problem

$$\begin{cases} D^\sigma u(t) + f(t, u(t)) = 0, & t \in [0, 1], \\ u^{(i)}(0) = 0, & i = 0, 1, 2, \dots, n - 2, \\ u(1) = \sum_{i=1}^{m-2} \beta_i \int_0^{\eta_i} u(s) ds + \sum_{i=1}^{m-2} \gamma_i u(\eta_i), \end{cases}$$

where D^σ represents the standard Riemann-Liouville fractional derivative of order σ satisfying $n - 1 < \sigma \leq n$ with $n \geq 3$. The authors used Krasnoselkii's fixed point theorem, Schauder type fixed point theorem, Banach's contraction mapping principle and nonlinear alternative for single-valued maps to obtain the existence results.

Inspired from the above works, in this paper, we establish the existence and multiplicity of positive solutions of the boundary value problem (1.1). Our paper is organized as follows. After this section, some definitions and lemmas will be established in Section 2. In Section 3, we give our main results in Theorems 2.10 and 2.11. Finally, in Section 4, as applications, some examples are presented to illustrate our main results.

2. PRELIMINARIES

In this section, we introduce some notations and definitions of fractional calculus, which can be found in [16, 24, 28]. We also state two fixed-point theorems due to Guo-Krasnosel'skii and Leggett-Williams.

Definition 2.1. *The Riemann-Liouville fractional integral of order $\alpha > 0$ for a function $f : (0, +\infty) \rightarrow \mathbb{R}$ is defined as*

$$I_{0+}^\alpha f(t) = \frac{1}{\Gamma(\alpha)} \int_0^t (t-s)^{\alpha-1} f(s) ds,$$

provided the right side is pointwise defined on $(0, +\infty)$ where $\Gamma(\cdot)$ is the Gamma function.

Definition 2.2. *The Riemann-Liouville fractional derivative order $\alpha > 0$ of a continuous function $u : (0, \infty) \rightarrow \mathbb{R}$ is defined by*

$$D_{0+}^\alpha u(t) = \frac{1}{\Gamma(n-\alpha)} \left(\frac{d}{dt} \right)^n \int_0^t (t-s)^{n-\alpha-1} u(s) ds,$$

where $n = [\alpha] + 1$, $[\alpha]$ denotes the integer part of number α , provided that the right side is pointwise defined on $(0, \infty)$.

Lemma 2.3. (i) If $u \in L^p(0, 1)$, $1 \leq p \leq +\infty$, $\beta > \alpha > 0$, then $I_{0+}^\alpha I_{0+}^\beta u(t) = I_{0+}^{\alpha+\beta} u(t)$.

(ii) If $\alpha > 0$ and $\gamma \in (-1, +\infty)$, then $I_{0+}^\alpha t^\gamma = \frac{\Gamma(\gamma+1)}{\Gamma(\alpha+\gamma+1)} t^{\alpha+\gamma}$.

Lemma 2.4. Let $\alpha > 0$ and for any $y \in L^1(0, 1)$. Then, the general solution of the fractional differential equation $D_{0+}^\alpha u(t) + y(t) = 0$, $0 < t < 1$ is given by

$$u(t) = -\frac{1}{\Gamma(\alpha)} \int_0^t (t-s)^{\alpha-1} y(s) ds + c_1 t^{\alpha-1} + c_2 t^{\alpha-2} + \dots + c_n t^{\alpha-n},$$

where c_0, c_1, \dots, c_{n-1} are real constants and $n = [\alpha] + 1$.

Definition 2.5. Let E be a real Banach space. A nonempty convex closed set $K \subset E$ is said to be a cone provided that

- (i) $au \in K$ for all $u \in K$ and all $a \geq 0$, and
- (ii) $u, -u \in K$ implies $u = 0$.

Definition 2.6. The map α is defined as a nonnegative continuous concave functional on a cone K of a real Banach space E provided that $\alpha : K \rightarrow [0, +\infty)$ is continuous and

$$\alpha(tx + (1-t)y) \geq t\alpha(x) + (1-t)\alpha(y)$$

for all $x, y \in K$ and $0 \leq t \leq 1$.

Lemma 2.7. Let $\Delta = 1 - \Gamma(q) \sum_{i=1}^m \frac{\alpha_i \eta^{p_i+q-1}}{\Gamma(p_i+q)} - \sum_{i=1}^m \beta_i \xi_i^{q-1} > 0$, $\alpha_i, \beta_i \geq 0$, $p_i > 0$, $i = 1, 2, \dots, m$, and $h \in C[0, 1]$. The unique the solution $u \in AC[0, 1]$ of the boundary value problem

$$D^q u(t) + h(t) = 0, \quad t \in (0, 1), \quad q \in (1, 2] \quad (2.1)$$

$$u(0) = 0, \quad u(1) = \sum_{i=1}^m \alpha_i (I^{p_i} u)(\eta) + \sum_{i=1}^m \beta_i u(\xi_i) \quad (2.2)$$

is given by

$$u(t) = \int_0^1 G(t, s) h(s) ds, \quad (2.3)$$

where $G(t, s)$ is the Green's function given by

$$G(t, s) = g(t, s) + \frac{t^{q-1}}{\Delta} \sum_{i=1}^m \frac{\alpha_i}{\Gamma(p_i+q)} g_i(\eta, s) + \frac{t^{q-1}}{\Delta} \sum_{i=1}^m \beta_i g(\xi_i, s) \quad (2.4)$$

where

$$g(t, s) = \frac{1}{\Gamma(q)} \begin{cases} t^{q-1} (1-s)^{q-1} - (t-s)^{q-1}, & 0 \leq s \leq t \leq 1, \\ t^{q-1} (1-s)^{q-1}, & 0 \leq t \leq s \leq 1, \end{cases} \quad (2.5)$$

and

$$g_i(\eta, s) = \begin{cases} \eta^{p_i+q-1} (1-s)^{q-1} - (\eta-s)^{p_i+q-1}, & 0 \leq s \leq \eta < 1, \\ \eta^{p_i+q-1} (1-s)^{q-1}, & 0 < \eta \leq s \leq 1, \end{cases} \quad (2.6)$$

Proof. By Lemma 2.4 the general solution for the above equation (2.1) is

$$u(t) = -\frac{1}{\Gamma(q)} \int_0^t (t-s)^{q-1} h(s) ds + c_1 t^{q-1} + c_2 t^{q-2},$$

where $c_1, c_2 \in \mathbb{R}$. The first condition of (2.2) implies that $c_2 = 0$. Thus

$$u(t) = -\frac{1}{\Gamma(q)} \int_0^t (t-s)^{q-1} h(s) ds + c_1 t^{q-1}. \quad (2.7)$$

Taking the Riemann-Liouville fractional integral of order $p_i > 0$ for (2.7) and using Lemma 2.3 we get that

$$\begin{aligned} (I^{p_i} u)(t) &= \int_0^t \frac{(t-s)^{p_i-1}}{\Gamma(p_i)} \left(c_1 s^{q-1} - \int_0^s \frac{(s-r)^{q-1}}{\Gamma(q)} dr \right) h(s) ds \\ &= c_1 \int_0^t \frac{(t-s)^{p_i-1} s^{q-1}}{\Gamma(p_i)} ds - \int_0^t \frac{(t-s)^{p_i-1}}{\Gamma(p_i)} \int_0^s \frac{(s-r)^{q-1}}{\Gamma(q)} h(r) ds dr \\ &= c_1 \frac{t^{p_i+q-1} \Gamma(q)}{\Gamma(p_i+q)} - \frac{1}{\Gamma(p_i+q)} \int_0^t (t-s)^{p_i+q-1} h(s) ds. \end{aligned}$$

The second condition of (2.2) yields

$$\begin{aligned} c_1 - \frac{1}{\Gamma(q)} \int_0^1 (1-s)^{q-1} h(s) ds &= c_1 \sum_{i=1}^m \frac{\alpha_i \eta^{p_i+q-1} \Gamma(q)}{\Gamma(p_i+q)} \\ &\quad - \sum_{i=1}^m \frac{\alpha_i}{\Gamma(p_i+q)} \int_0^\eta (\eta-s)^{p_i+q-1} h(s) ds \\ &\quad + c_1 \sum_{i=1}^m \beta_i \xi_i^{q-1} - \frac{1}{\Gamma(q)} \sum_{i=1}^m \beta_i \int_0^{\xi_i} (\xi_i-s) h(s) ds. \end{aligned}$$

Then, we have that

$$\begin{aligned} c_1 &= \frac{1}{\Delta} \left\{ \int_0^1 \frac{(1-s)^{q-1}}{\Gamma(q)} h(s) ds - \sum_{i=1}^m \frac{\alpha_i}{\Gamma(p_i+q)} \int_0^\eta (\eta-s)^{p_i+q-1} h(s) ds \right. \\ &\quad \left. - \frac{1}{\Gamma(q)} \sum_{i=1}^m \beta_i \int_0^{\xi_i} (\xi_i-s)^{q-1} h(s) ds \right\}. \end{aligned}$$

Hence, the solution is

$$\begin{aligned}
 u(t) &= -\frac{1}{\Gamma(q)} \int_0^t (t-s)^{q-1} h(s) ds + \frac{t^{q-1}}{\Delta\Gamma(q)} \int_0^1 (1-s)^{q-1} h(s) ds \\
 &\quad - \frac{t^{q-1}}{\Delta} \sum_{i=1}^m \frac{\alpha_i}{\Gamma(p_i+q)} \int_0^n (\eta-s)^{p_i+q-1} h(s) ds - \frac{t^{q-1}}{\Delta\Gamma(q)} \sum_{i=1}^m \beta_i \int_0^{\xi_i} (\xi_i-s)^{q-1} h(s) ds \\
 &= -\frac{1}{\Gamma(q)} \int_0^t (t-s)^{q-1} h(s) ds + \frac{t^{q-1}}{\Gamma(q)} \int_0^1 (1-s)^{q-1} h(s) ds \\
 &\quad + \frac{t^{q-1}}{\Delta} \left\{ \sum_{i=1}^m \frac{\alpha_i \eta^{p_i+q-1}}{\Gamma(p_i+q)} + \frac{1}{\Gamma(q)} \sum_{i=1}^m \beta_i \xi_i^{q-1} \right\} \int_0^1 (1-s)^{q-1} h(s) ds \\
 &\quad - \frac{t^{q-1}}{\Delta} \sum_{i=1}^m \frac{\alpha_i}{\Gamma(p_i+q)} \int_0^n (\eta-s)^{p_i+q-1} h(s) ds - \frac{t^{q-1}}{\Delta\Gamma(q)} \sum_{i=1}^m \beta_i \int_0^{\xi_i} (\xi_i-s)^{q-1} h(s) ds \\
 &= \int_0^1 g(t,s) h(s) ds + \frac{t^{q-1}}{\Delta} \sum_{i=1}^m \frac{\alpha_i}{\Gamma(p_i+q)} \int_0^1 g_i(\eta,s) h(s) ds \\
 &\quad + \frac{t^{q-1}}{\Delta} \sum_{i=1}^m \beta_i \int_0^1 g(\xi_i,s) h(s) ds \\
 &= \int_0^1 G(t,s) h(s) ds.
 \end{aligned}$$

□

Lemma 2.8. *The Green's function $G(t,s)$ has the following properties:*

- (P₁) $G(t,s)$ is continuous on $[0,1] \times [0,1]$.
- (P₂) $G(t,s) \geq 0$ for all $0 \leq s, t \leq 1$.
- (P₃) $G(t,s) \leq \max_{0 \leq t \leq 1} G(t,s) \leq g(s,s) \left(1 + \frac{\sum_{i=1}^m \beta_i}{\Delta} \right) + \sum_{i=1}^m \frac{\alpha_i}{\Delta\Gamma(p_i+q)} g_i(\eta,s)$
- (P₄) $\int_0^1 \max_{0 \leq t \leq 1} G(t,s) ds \leq \left(1 + \frac{\sum_{i=1}^m \beta_i}{\Delta} \right) \frac{\Gamma(q)}{\Gamma(2q)} + \sum_{i=1}^m \frac{\alpha_i \eta^{p_i+q-1}}{\Delta\Gamma(p_i+q)} \left(\frac{p_i+q(1-\eta)}{q(p_i+q)} \right)$
- (P₅) $\min_{\eta \leq t \leq 1} G(t,s) \geq \sum_{i=1}^m \frac{\alpha_i \eta^{q-1}}{\Delta\Gamma(p_i+q)} g_i(\eta,s) + (q-1) \sum_{i=1}^m \frac{\beta_i (\xi_i^{q-1} - \xi_i^q) \eta^{q-1}}{\Delta} s g(s,s)$ for $s \in [0,1]$.

Proof. It is easy to check that (P₁) holds. To prove (P₂), we will show that $g(t,s) \geq 0$ and $g_i(\eta,s) \geq 0$, $i = 1, 2, \dots, m$, for all $0 \leq s, t \leq 1$. For $t \leq s$, it is clear that $G(t,s) \geq 0$, we only need to prove the case $s \leq t$.

Then

$$\begin{aligned}
 g(t,s) &= \frac{1}{\Gamma(q)} \left[t^{q-1} (1-s)^{q-1} - (t-s)^{q-1} \right] \\
 &= \frac{1}{\Gamma(q)} \left[(t-ts)^{q-1} - (t-s)^{q-1} \right] \\
 &\geq \frac{1}{\Gamma(q)} \left[(t-s)^{q-1} - (t-s)^{q-1} \right] = 0.
 \end{aligned}$$

For $0 \leq s \leq \eta < 1$, we have

$$\begin{aligned}
g_i(\eta, s) &= \eta^{p_i+q-1} (1-s)^{q-1} - (\eta-s)^{p_i+q-1} \\
&= \eta^{p_i} (\eta-\eta s)^{q-1} - (\eta-s)^{p_i+q-1} \\
&\geq \eta^{p_i} (\eta-s)^{q-1} - (\eta-s)^{p_i+q-1} \\
&= (\eta-s)^{q-1} (\eta^{p_i} - (\eta-s)^{p_i}) \\
&\geq 0.
\end{aligned}$$

When $0 < \eta \leq s \leq 1$, $g_i(\eta, s) = \eta^{p_i+q-1} (1-s)^{q-1} \geq 0$. Therefore, $g_i(\eta, s) \geq 0$, $i = 1, 2, \dots, m$ for all $0 \leq s \leq 1$

Now, we prove (P_3) . For a given $s \in [0, 1]$, when $0 \leq s \leq t \leq 1$

$$\Gamma(q) g(t, s) = t^{q-1} (1-s)^{q-1} - (t-s)^{q-1}$$

and thus

$$\begin{aligned}
\Gamma(q) \frac{\partial}{\partial t} g(t, s) &= (q-1) t^{q-2} (1-s)^{q-1} - (q-1) (t-s)^{q-2} \\
&= (q-1) (t-ts)^{q-2} (1-s) - (q-1) (t-s)^{q-2} \\
&\leq (q-1) (t-s)^{q-2} (1-s) - (q-1) (t-s)^{q-2} \\
&= -s(q-1) (t-s)^{q-2}.
\end{aligned}$$

Hence, $g(t, s)$ is decreasing with respect to t . Then we have $g(t, s) \leq g(s, s)$ for $0 \leq s \leq t \leq 1$. For $0 \leq t \leq s \leq 1$

$$\Gamma(q) \frac{\partial}{\partial t} g(t, s) = (q-1) t^{q-2} (1-s)^{q-1} \geq 0,$$

which means that $g(t, s)$ is increasing with respect to t . Thus $g(t, s) \leq g(s, s)$ for $0 \leq t \leq s \leq 1$. Therefore $g(t, s) \leq g(s, s)$ for $0 \leq s, t \leq 1$

From the above analysis, we have for $0 \leq s \leq 1$ that

$$\begin{aligned}
G(t, s) &\leq \max_{0 \leq t \leq 1} G(t, s) = \max_{0 \leq t \leq 1} (g(t, s) \\
&\quad + \frac{t^{q-1}}{\Delta} \sum_{i=1}^m \frac{\alpha_i}{\Gamma(p_i+q)} g_i(\eta, s) + \frac{t^{q-1}}{\Delta} \sum_{i=1}^m \beta_i g(\xi_i, s)) \\
&\leq g(s, s) \left(1 + \frac{\sum_{i=1}^m \beta_i}{\Delta} \right) + \sum_{i=1}^m \frac{\alpha_i}{\Delta \Gamma(p_i+q)} g_i(\eta, s).
\end{aligned}$$

To prove (P_4) , by direct integration, we have

$$\begin{aligned}
 \int_0^1 \max_{0 \leq t \leq 1} G(t, s) ds &\leq \int_0^1 \left[g(s, s) \left(1 + \frac{\sum_{i=1}^m \beta_i}{\Delta} \right) + \sum_{i=1}^m \frac{\alpha_i}{\Delta \Gamma(p_i + q)} g_i(\eta, s) \right] ds \\
 &= \left(1 + \frac{\sum_{i=1}^m \beta_i}{\Delta} \right) \int_0^1 \frac{s^{q-1} (1-s)^{q-1}}{\Gamma(q)} ds \\
 &\quad + \sum_{i=1}^m \frac{\alpha_i}{\Delta \Gamma(p_i + q)} \left(\int_\eta^1 \eta^{p_i+q-1} (1-s)^{q-1} \right) ds \\
 &\quad + \sum_{i=1}^m \frac{\alpha_i}{\Delta \Gamma(p_i + q)} \left(\int_0^\eta \left[\eta^{p_i+q-1} (1-s)^{q-1} - (\eta-s)^{p_i+q-1} \right] ds \right) \\
 &= \left(1 + \frac{\sum_{i=1}^m \beta_i}{\Delta} \right) \frac{\Gamma(q)}{\Gamma(2q)} + \sum_{i=1}^m \frac{\alpha_i \eta^{p_i+q-1}}{\Delta \Gamma(p_i + q)} \left(\frac{p_i + q(1-\eta)}{q(p_i + q)} \right).
 \end{aligned}$$

Now, we shall prove (P_5) .

Firstly, let $k_1(\xi_i, s) = \frac{g(\xi_i, s)}{g(s, s)}$ for $0 < s < \xi_i < 1$, $i = 1, 2, \dots, m$, then we get

$$k_1(\xi_i, s) = \frac{(\xi_i(1-s))^{q-1} - (\xi_i - s)^{q-1}}{s^{q-1}(1-s)^{q-1}} = \frac{(q-1) \int_{\xi_i - s}^{\xi_i(1-s)} x^{q-2} dx}{s^{q-1}(1-s)^{q-1}}$$

Since the function $x \mapsto x^{q-2}$ is continuous and decreasing on $[\xi_i - s, \xi_i(1-s)]$, we have

$$\begin{aligned}
 k_1(\xi_i, s) &\geq \frac{(q-1) (\xi_i(1-s))^{q-2} [\xi_i(1-s) - (\xi_i - s)]}{s^{q-1}(1-s)^{q-1}} \\
 &= \frac{(q-1) \xi_i^{q-2} (1-s)^{q-2} s(1-\xi_i)}{s^{q-1}(1-s)^{q-1}} \\
 &\geq (q-1) \xi_i^{q-1} (1-\xi_i) s.
 \end{aligned}$$

Let $k_2(\xi_i, s) = \frac{g(\xi_i, s)}{g(s, s)}$ for $0 < \xi_i \leq s < 1$, $i = 1, 2, \dots, m$, then we get

$$k_2(\xi_i, s) = \frac{\xi_i^{q-1}}{s^{q-1}} \geq \frac{\xi_i^{q-1}}{s^{q-2}} = \xi_i^{q-1} s^{2-q} \geq (q-1) \xi_i^{q-1} (1-\xi_i) s$$

Therefore, we have

$$g(\xi_i, s) \geq (q-1) s g(s, s) \left(\xi_i^{q-1} - \xi_i^q \right) \quad \text{for } 0 < s, \xi_i < 1 \quad (2.8)$$

Furthermore, the inequality in (2.8) is satisfied for $s \in \{0, 1\}$. Hence

$$g(\xi_i, s) \geq (q-1) s g(s, s) \left(\xi_i^{q-1} - \xi_i^q \right) \quad \text{for } 0 \leq s, \xi_i \leq 1. \quad (2.9)$$

Secondly, from $g(t, s) \geq 0$, $g_i(\eta, s) \geq 0$, $i = 1, 2, \dots, m$ and from (2.9), we have

$$\begin{aligned}
\min_{\eta \leq t \leq 1} G(t, s) &= \min_{\eta \leq t \leq 1} \left(g(t, s) + \frac{t^{q-1}}{\Delta} \sum_{i=1}^m \frac{\alpha_i}{\Gamma(p_i + q)} g_i(\eta, s) \right. \\
&\quad \left. + \frac{t^{q-1}}{\Delta} \sum_{i=1}^m \beta_i g(\xi_i, s) \right) \\
&\geq \min_{\eta \leq t \leq 1} g(t, s) + \min_{\eta \leq t \leq 1} \frac{t^{q-1}}{\Delta} \sum_{i=1}^m \frac{\alpha_i}{\Gamma(p_i + q)} g_i(\eta, s) \\
&\quad + \min_{\eta \leq t \leq 1} \frac{t^{q-1}}{\Delta} \sum_{i=1}^m \beta_i g(\xi_i, s) \\
&\geq \sum_{i=1}^m \frac{\alpha_i \eta^{q-1}}{\Delta \Gamma(p_i + q)} g_i(\eta, s) + (q-1) \sum_{i=1}^m \frac{\beta_i (\xi_i^{q-1} - \xi_i^q) \eta^{q-1}}{\Delta} s g(s, s)
\end{aligned}$$

for $0 \leq s \leq 1$. This completes the proof. \square

Let $E = C([0, 1], \mathbb{R})$ be the Banach space of all continuous functions defined on $[0, 1]$ that are mapped into \mathbb{R} with the norm defined as $\|u\| = \sup_{t \in [0, 1]} |u(t)|$. If $u \in E$ satisfies the problem (1.1) and $u(t) \geq 0$ for any $t \in [0, 1]$, then u is called a nonnegative solution of the problem (1.1). If u is a nonnegative solution of the problem (1.1) with $\|u\| > 0$, then u is called a positive solution of the problem (1.1). Define the cone $\mathcal{K} \in E$ by

$$\mathcal{K} = \{u \in E : u(t) \geq 0\},$$

and the operator $A : \mathcal{K} \rightarrow E$ by

$$Au(t) := \int_0^1 G(t, s) f(s, u(s)) ds. \quad (2.10)$$

In view of Lemma 2.7, the nonnegative solutions of problem (1.1) are given by the operator equation $u(t) = Au(t)$

Lemma 2.9. *Suppose that $f : [0, 1] \times [0, \infty) \rightarrow [0, \infty)$ is continuous. The operator $A : \mathcal{K} \rightarrow \mathcal{K}$ is completely continuous.*

Proof. Since $G(t, s) \geq 0$ for $s, t \in [0, 1]$, we have $Au(t) \geq 0$ for all $u \in \mathcal{K}$. Therefore, $A : \mathcal{K} \rightarrow \mathcal{K}$.

For a constant $R > 0$, we define $\Omega = \{u \in \mathcal{K} : \|u\| < R\}$.

Let

$$L = \max_{0 \leq t \leq 1, 0 \leq u \leq R} |f(t, u)|. \quad (2.11)$$

Then, for $u \in \Omega$, from Lemma 2.8, we have

$$\begin{aligned}
 |Au(t)| &= \left| \int_0^1 G(t,s) f(s, u(s)) ds \right| \\
 &\leq L \int_0^1 G(t,s) ds \\
 &\leq L \int_0^1 \left(g(s,s) \left(1 + \frac{\sum_{i=1}^m \beta_i}{\Delta} \right) + \sum_{i=1}^m \frac{\alpha_i}{\Delta \Gamma(p_i + q)} g_i(\eta, s) \right) ds \quad (2.12) \\
 &\leq \left(1 + \frac{\sum_{i=1}^m \beta_i}{\Delta} \right) \frac{\Gamma(q)}{\Gamma(2q)} + \sum_{i=1}^m \frac{\alpha_i \eta^{p_i + q - 1}}{\Delta \Gamma(p_i + q)} \left(\frac{p_i + q(1 - \eta)}{q(p_i + q)} \right).
 \end{aligned}$$

Hence, $\|Au\| \leq M$, and so $A(\Omega)$ is uniformly bounded. Now, we shall show that $A(\Omega)$ is equicontinuous. For $u \in \Omega$, $t_1, t_2 \in [0, 1]$, $t_1 < t_2$, we have

$$|Au(t_2) - Au(t_1)| \leq L \int_0^1 |G(t_2, s) - G(t_1, s)| ds,$$

where L is defined by (2.11). Since $G(t, s)$ is continuous on $[0, 1] \times [0, 1]$, therefore $G(t, s)$ is uniformly continuous on $[0, 1] \times [0, 1]$. Hence, for any $\epsilon > 0$, there exists a positive constant

$$\delta = \frac{1}{2} \left[\frac{\epsilon \Gamma(q)}{L} \left(\frac{1}{\frac{1}{q} + \sum_{i=1}^m \frac{\alpha_i \eta^{p_i + q - 1}}{\Delta \Gamma(p_i + q)} \left(\frac{p_i + q(1 - \eta)}{q(p_i + q)} \right) + \frac{\Gamma(q)}{\Gamma(2q)} \frac{\sum_{i=1}^m \beta_i}{\Delta}} \right) \right]$$

whenever $|t_2 - t_1| < \delta$, we have the following two cases.

Case 1. $\delta \leq t_1 < t_2 < 1$

Therefore,

$$\begin{aligned}
|Au(t_2) - Au(t_1)| &\leq L \int_0^1 |G(t_2, s) - G(t_1, s)| ds \\
&= L \left[\int_0^{t_1} |G(t_2, s) - G(t_1, s)| ds + \int_{t_1}^{t_2} |G(t_2, s) - G(t_1, s)| ds \right. \\
&\quad \left. + \int_{t_2}^1 |G(t_2, s) - G(t_1, s)| ds \right] \\
&\leq \frac{(t_2^{q-1} - t_1^{q-1})L}{\Gamma(q)} \int_0^1 (1-s)^{q-1} ds + \frac{(t_2^{q-1} - t_1^{q-1})L}{\Delta} \int_0^1 \sum_{i=1}^m \frac{\alpha_i}{\Gamma(p_i + q)} g_i(\eta, s) ds \\
&\quad + \frac{(t_2^{q-1} - t_1^{q-1})L}{\Delta} \sum_{i=1}^m \beta_i \int_0^1 g(s, s) ds \\
&= (t_2^{q-1} - t_1^{q-1})L \left[\frac{1}{q\Gamma(q)} + \sum_{i=1}^m \frac{\alpha_i \eta^{p_i+q-1}}{\Delta \Gamma(p_i + q)} \left(\frac{p_i + q(1-n)}{q(p_i + q)} \right) + \frac{\Gamma(q)}{\Gamma(2q)} \frac{\sum_{i=1}^m \beta_i}{\Delta} \right] \\
&= \frac{(t_2^{q-1} - t_1^{q-1})L}{\Gamma(q)} \left[\frac{1}{q} + \sum_{i=1}^m \frac{\alpha_i \eta^{p_i+q-1}}{\Delta \Gamma(p_i + q)} \left(\frac{p_i + q(1-n)}{q(p_i + q)} \right) + \frac{\Gamma(q)}{\Gamma(2q)} \frac{\sum_{i=1}^m \beta_i}{\Delta} \right] \\
&\leq \frac{(q-1)\delta^{q-1}L}{\Gamma(q)} \left[\frac{1}{q} + \sum_{i=1}^m \frac{\alpha_i \eta^{p_i+q-1}}{\Delta \Gamma(p_i + q)} \left(\frac{p_i + q(1-n)}{q(p_i + q)} \right) + \frac{\Gamma(q)}{\Gamma(2q)} \frac{\sum_{i=1}^m \beta_i}{\Delta} \right] \\
&< \epsilon.
\end{aligned}$$

Case 2. $0 \leq t_1 < 1, t_2 < 2\delta$

Hence

$$\begin{aligned}
|Au(t_2) - Au(t_1)| &\leq L \int_0^1 |G(t_2, s) - G(t_1, s)| ds \\
&< \frac{(t_2^{q-1} - t_1^{q-1})L}{\Gamma(q)} \left[\frac{1}{q} + \sum_{i=1}^m \frac{\alpha_i \eta^{p_i+q-1}}{\Delta \Gamma(p_i + q)} \left(\frac{p_i + q(1-n)}{q(p_i + q)} \right) + \frac{\Gamma(q)}{\Gamma(2q)} \frac{\sum_{i=1}^m \beta_i}{\Delta} \right] \\
&\leq \frac{t_2^{q-1}L}{\Gamma(q)} \left[\frac{1}{q} + \sum_{i=1}^m \frac{\alpha_i \eta^{p_i+q-1}}{\Delta \Gamma(p_i + q)} \left(\frac{p_i + q(1-n)}{q(p_i + q)} \right) + \frac{\Gamma(q)}{\Gamma(2q)} \frac{\sum_{i=1}^m \beta_i}{\Delta} \right] \\
&< \frac{(2\delta)^{q-1}L}{\Gamma(q)} \left[\frac{1}{q} + \sum_{i=1}^m \frac{\alpha_i \eta^{p_i+q-1}}{\Delta \Gamma(p_i + q)} \left(\frac{p_i + q(1-n)}{q(p_i + q)} \right) + \frac{\Gamma(q)}{\Gamma(2q)} \frac{\sum_{i=1}^m \beta_i}{\Delta} \right] \\
&= \epsilon.
\end{aligned}$$

Thus, $A(\Omega)$ is equicontinuous. In view of the Arzela-Ascoli theorem, we have that $\overline{A(\Omega)}$ is compact, which means $A : \mathcal{K} \rightarrow \mathcal{K}$ is a completely continuous operator. This completes the proof. \square

Theorem 2.10. [10] *Let E be a Banach space, and let $\mathcal{K} \in E$ be a cone. Assume that Ω_1, Ω_2 are open subsets of E with $0 \in \Omega_1, \overline{\Omega_1} \subset \Omega_2$, and let $T : \mathcal{K} \cap (\Omega_2 \setminus \Omega_1) \rightarrow \mathcal{K}$ be a completely continuous operator such that:*

(i) $\|Tu\| \geq \|u\|$, $u \in \mathcal{K} \cap \partial\Omega_1$, and $\|Tu\| \leq \|u\|$, $u \in \mathcal{K} \cap \partial\Omega_2$; or

(ii) $\|Tu\| \leq \|u\|$, $u \in \mathcal{K} \cap \partial\Omega_1$, and $\|Tu\| \geq \|u\|$, $u \in \mathcal{K} \cap \partial\Omega_2$.

Then T has a fixed point $\mathcal{K} \cap (\overline{\Omega_2} \setminus \Omega_1)$.

Theorem 2.11. [17] *Let \mathcal{K} be a cone in the real Banach space E and $c > 0$ be a constant. Assume that there exists a concave nonnegative continuous functional θ on \mathcal{K} with $\theta(u) \leq \|u\|$ for all $u \in \overline{\mathcal{K}_c}$. Let $A : \overline{\mathcal{K}_c} \rightarrow \overline{\mathcal{K}_c}$ be a completely continuous operator. Suppose that there exist constants $0 < a < b < d \leq c$ such that the following conditions hold:*

(i) $\{u \in \mathcal{K}(\theta, b, d) : \theta(u) > b\} \neq \emptyset$ and $\theta(Au) > b$ for $u \in \mathcal{K}(\theta, b, d)$;

(ii) $\|Au\| < a$ for $\|u\| \leq a$;

(iii) $\theta(Au) > b$ for $u \in \mathcal{K}(\theta, b, c)$ with $\|Au\| > d$.

Then A has at least three fixed points u_1 , u_2 and u_3 in $\overline{\mathcal{K}_c}$ such that $\|u_1\| < a$, $b < \theta(u_2)$, $a < \|u_3\|$ with $\theta(u_3) < b$.

Remark 2.12. *If there holds $d = c$, then condition (i) implies condition (iii) of Theorem 2.11.*

3. MAIN RESULTS

In this section, in order to establish some results of existence and multiplicity of positive solutions for BVP (1.1), we will impose growth conditions on f which allow us to apply Theorems 2.10 and 2.11.

For convenience, we denote

$$\begin{aligned} \Lambda_1 &= \sum_{i=1}^m \frac{\alpha_i \eta^{p_i+2(q-1)}}{\Delta \Gamma(p_i+q)} \left(\frac{p_i+q(1-\eta)}{q(p_i+q)} \right) + (q-1) \sum_{i=1}^m \frac{\beta_i \left(\xi_i^{q-1} - \xi_i^q \right) \eta^{q-1}}{\Delta} \times \frac{\Gamma(q+1)}{\Gamma(2q+1)} \\ \Lambda_2 &= \left(1 + \frac{\sum_{i=1}^m \beta_i}{\Delta} \right) \frac{\Gamma(q)}{\Gamma(2q)} + \sum_{i=1}^m \frac{\alpha_i \eta^{p_i+q-1}}{\Delta \Gamma(p_i+q)} \left(\frac{p_i+q(1-\eta)}{q(p_i+q)} \right) \\ \Lambda_3 &= \sum_{i=1}^m \frac{\alpha_i \eta^{p_i+2(q-1)} (1-\eta)^q}{\Delta \Gamma(p_i+q) q} + (q-1) \sum_{i=1}^m \frac{\beta_i \left(\xi_i^{q-1} - \xi_i^q \right) \eta^{q-1} (1-\eta)^{2q} \Gamma(q+1)}{\Delta \Gamma(2q+1)} \end{aligned}$$

Theorem 3.1. *Let $f : [0, 1] \times [0, \infty) \rightarrow [0, \infty)$ be a continuous function. Assume that there exist constants $r_2 > r_1 > 0$, $M_1 \in (\Lambda_1^{-1}, \infty)$ and $M_2 \in (0, \Lambda_2^{-1})$ such that:*

(H₁) $f(t, u) \geq M_1 r_1$, for $(t, u) \in [0, 1] \times [0, r_1]$;

(H₂) $f(t, u) \leq M_2 r_2$, for $(t, u) \in [0, 1] \times [0, r_2]$.

Then boundary value problem (1.1) has at least one positive solution u such that $r_1 \leq \|u\| \leq r_2$.

Proof. From Lemma 2.9, the operator $A : \mathcal{K} \rightarrow \mathcal{K}$ is completely continuous. We divide the rest of the proof into two steps.

Step 1. Let $\Omega_1 = \{u \in E : \|u\| < r_1\}$, then for any $u \in \mathcal{K} \cap \Omega_1$, we have $0 \leq u(t) \leq r_1$ for all $t \in [0, 1]$. From (H_1) , it follows for $t \in [\eta, 1]$ that

$$\begin{aligned}
(Au)(t) &= \int_0^1 G(t, s) f(s, u(s)) ds \\
&\geq \int_0^1 \min_{\eta \leq t \leq 1} G(t, s) f(s, u(s)) ds \\
&\geq M_1 r_1 \left\{ \sum_{i=1}^m \frac{\alpha_i \eta^{q-1}}{\Delta \Gamma(p_i + q)} \int_0^1 g_i(\eta, s) ds + (q-1) \sum_{i=1}^m \frac{\beta_i (\xi_i^{q-1} - \xi_i)}{\Delta} \int_0^1 sg(s, s) ds \right\} \\
&= M_1 r_1 \left\{ \sum_{i=1}^m \frac{\alpha_i \eta^{q-1}}{\Delta \Gamma(p_i + q)} \left(\int_\eta^1 \eta^{p_i+q-1} (1-s)^{q-1} ds \right. \right. \\
&\quad \left. \left. + \int_0^\eta \left[\eta^{p_i+q-1} (1-s)^{q-1} - (\eta-s)^{p_i+q-1} \right] ds \right) \right. \\
&\quad \left. + (q-1) \sum_{i=1}^m \frac{\beta_i (\xi_i^{q-1} - \xi_i^q) \eta^{q-1}}{\Delta} \times \frac{\Gamma(q+1)}{\Gamma(2q+1)} \right\} \\
&= M_1 r_1 \left\{ \sum_{i=1}^m \frac{\alpha_i \eta^{p_i+2(q-1)}}{\Delta \Gamma(p_i + q)} \left(\frac{p_i + q(1-\eta)}{q(p_i + q)} \right) + (q-1) \sum_{i=1}^m \frac{\beta_i (\xi_i^{q-1} - \xi_i^q) \eta^{q-1}}{\Delta} \times \frac{\Gamma(q+1)}{\Gamma(2q+1)} \right\} \\
&\geq r_1 = \|u\|,
\end{aligned}$$

which means that

$$\|Au\| \geq \|u\| \quad \text{for } u \in \mathcal{K} \cap \partial\Omega_1. \quad (3.1)$$

Step 2. Let $\Omega_2 = \{u \in E : \|u\| < r_2\}$, then for any $u \in \mathcal{K} \cap \partial\Omega_2$, we have $0 \leq u(t) \leq r_2$ for all $t \in [0, 1]$. It follows from (H_2) that for $t \in [0, 1]$,

$$\begin{aligned}
(Au)(t) &= \int_0^1 G(t, s) f(s, u(s)) ds \\
&\leq M_2 r_2 \left\{ \left(1 + \frac{\sum_{i=1}^m \beta_i}{\Delta} \right) \frac{\Gamma(q)}{\Gamma(2q)} + \sum_{i=1}^m \frac{\alpha_i \eta^{p_i+q-1}}{\Delta \Gamma(p_i + q)} \left(\frac{p_i + q(1-\eta)}{q(p_i + q)} \right) \right\} \\
&\leq r_2 = \|u\|,
\end{aligned}$$

which means that

$$\|Au\| \leq \|u\| \quad \text{for any } u \in \mathcal{K} \cap \partial\Omega_2. \quad (3.2)$$

By (i) of Theorem 2.10, we get that A has a fixed point u in \mathcal{K} with $r_1 \leq \|u\| \leq r_2$, which is also a positive solution of boundary value problem (1.1). \square

Theorem 3.2. *Let $f : [0, 1] \times [0, \infty) \rightarrow [0, \infty)$ be a continuous function. Suppose that there exist constants $0 < a < b < c$ such that the following assumptions hold:*

(H_3) $f(t, u) < \Lambda_2^{-1}a$ for $(t, u) \in [0, 1] \times [0, a]$;

(H_4) $f(t, u) > \Lambda_3^{-1}b$ for $(t, u) \in [\eta, 1] \times [b, c]$;

(H_5) $f(t, u) \leq \Lambda_2^{-1}c$ for $(t, u) \in [0, 1] \times [0, c]$.

Then boundary value problem (1.1) has at least one nonnegative solution u_1 and two positive solutions u_2, u_3 in $\bar{\mathcal{K}}_c$ with

$$\|u_1\| < a, \quad b < \min_{\eta \leq t \leq 1} u_2(t) \text{ and } a < \|u_3\| \quad \text{with} \quad \min_{\eta \leq t \leq 1} u_3(t) < b.$$

Proof. We show that all the conditions of Theorem 2.11 are satisfied.

If $u \in \overline{\mathcal{K}}_c$, then $\|u\| \leq c$. Condition (H_5) implies $f(t, u(t)) \leq \Lambda_2^{-1}c$ for $t \in [0, 1]$. Consequently,

$$\begin{aligned} (Au)(t) &= \int_0^1 G(t, s) f(s, u(s)) ds \\ &\leq \Lambda_2^{-1}c \int_0^1 \left[\left(1 + \frac{\sum_{i=1}^m \beta_i}{\Delta}\right) g(s, s) + \sum_{i=1}^m \frac{\alpha_i \eta^{p_i+q-1}}{\Delta \Gamma(p_i+q)} \left(\frac{p_i+q(1-\eta)}{q(p_i+q)}\right) g_i(\eta, s) \right] ds \\ &= \Lambda_2^{-1}c \left\{ \left(1 + \frac{\sum_{i=1}^m \beta_i}{\Delta}\right) \frac{\Gamma(q)}{\Gamma(2q)} + \sum_{i=1}^m \frac{\alpha_i \eta^{p_i+q-1}}{\Delta \Gamma(p_i+q)} \left(\frac{p_i+q(1-\eta)}{q(p_i+q)}\right) \right\} \\ &= c, \end{aligned}$$

which implies $\|Au\| \leq c$. Hence, $A : \overline{\mathcal{K}}_c \rightarrow \overline{\mathcal{K}}_c$ is completely continuous.

If $u \in \overline{\mathcal{K}}_a$, then (H_3) yields

$$\begin{aligned} (Au)(t) &< \Lambda_2^{-1} \int_0^1 \left[\left(1 + \frac{\sum_{i=1}^m \beta_i}{\Delta}\right) g(s, s) + \sum_{i=1}^m \frac{\alpha_i}{\Delta \Gamma(p_i+q)} g_i(\eta, s) \right] ds \\ &= \Lambda_2^{-1}a \left\{ \left(1 + \frac{\sum_{i=1}^m \beta_i}{\Delta}\right) \frac{\Gamma(q)}{\Gamma(2q)} + \sum_{i=1}^m \frac{\alpha_i \eta^{p_i+q-1}}{\Delta \Gamma(p_i+q)} \left(\frac{p_i+q(1-\eta)}{q(p_i+q)}\right) \right\} \\ &= a. \end{aligned}$$

Thus $\|Au\| < a$. Therefore, condition (ii) of Theorem 2.11 holds.

Define a concave nonnegative continuous functional θ on \mathcal{K} by $\theta(u) = \min_{\eta \leq t \leq 1} |u(t)|$. To check condition (i) of Theorem 2.11, we choose $u(t) = \frac{b+c}{2}$ for $t \in [0, 1]$. It is easy to see that $u(t) \in \mathcal{K}(\theta, b, c)$ and $\theta(u) = \theta\left(\frac{b+c}{2}\right) > b$, which means that $\{\mathcal{K}(\theta, b, c) : \theta(u) > b\} \neq \emptyset$. Hence, if $u \in \mathcal{K}(\theta, b, c)$, then $b \leq u(t) \leq c$ for $t \in [\eta, 1]$. From assumption (H_4) , we have

$$\begin{aligned} \theta(Au) &= \min_{\eta \leq t \leq 1} |(Au)(t)| \\ &\geq \int_{\eta}^1 \min_{\eta \leq t \leq 1} G(t, s) f(s, u(s)) ds \\ &> \Lambda_3^{-1}b \left\{ \sum_{i=1}^m \frac{\alpha_i \eta^{q-1}}{\Delta \Gamma(p_i+q)} \int_{\eta}^1 g_i(\eta, s) ds + (q-1) \sum_{i=1}^m \frac{\beta_i (\xi_i^{q-1} - \xi_i) \eta^{q-1}}{\Delta} \int_{\eta}^1 g(s, s) ds \right\} \\ &= \Lambda_3^{-1}b \left\{ \sum_{i=1}^m \frac{\alpha_i \eta^{p_i+2(q-1)} (1-\eta)^q}{\Delta \Gamma(p_i+q) q} + (q-1) \sum_{i=1}^m \frac{\beta_i (\xi_i^q - \xi_i^q) \eta^{q-1} (1-\eta)^{2q} \Gamma(q+1)}{\Delta \Gamma(2q+1)} \right\} \\ &= b. \end{aligned}$$

Thus $\theta(Au) > b$ for all $u \in \mathcal{K}(\theta, b, c)$. This shows that condition (i) of Theorem 2.11 is also satisfied.

By Theorem 2.11 and Remark 2.12, boundary value problem (1.1) has at least one nonnegative solution u_1 and two positive solutions u_2, u_3 , which satisfy

$$\|u_1\| < a, \quad b < \min_{\eta \leq t \leq 1} |u_2(t)| \quad a < \|u_3\| \quad \text{with} \quad \min_{\eta \leq t \leq 1} |u(t)| < b.$$

The proof is complete. \square

4. EXAMPLES

4.1. Example. Consider the fractional differential equations with boundary value as follows:

$$\begin{cases} D^{\frac{3}{2}}u(t) + f(t, u(t)) = 0, & 0 < t < 1, \\ u(0) = 0 \\ u(1) = 2 \left(I^{\frac{3}{2}}u \right) \left(\frac{1}{4} \right) + \frac{1}{2} \left(I^{\frac{\pi}{4}}u \right) \left(\frac{1}{4} \right) + \frac{4}{5} \left(I^{\frac{2}{3}}u \right) \left(\frac{1}{4} \right) + \frac{3}{15}u \left(\frac{1}{3} \right) + \frac{3}{20}u \left(\frac{1}{4} \right) + \frac{1}{4}u \left(\frac{1}{5} \right), \end{cases} \quad (4.1)$$

where

$$f(t, u) \begin{cases} u(1 - u^2) + 4 \left(1 + \frac{3}{4}t \right), & 0 \leq t \leq 1; 0 \leq u \leq 13 \\ 4 \left(1 + \frac{3}{4}t \right) e^{1-u} + \sin^2(\pi(1-x)), & 0 \leq t \leq 1; 1 \leq u \leq 20 \end{cases}$$

Set $m = 3$, $\eta = \frac{1}{4}$, $q = \frac{3}{2}$, $\alpha_1 = 2$, $\alpha_2 = \frac{1}{2}$, $\alpha_3 = \frac{4}{5}$, $p_1 = \frac{3}{2}$, $p_2 = \frac{\pi}{4}$, $p_3 = \frac{2}{3}$, $\beta_1 = \frac{1}{4}$, $\beta_2 = \frac{3}{20}$, $\beta_3 = \frac{3}{5}$, $\xi_1 = \frac{1}{5}$, $\xi_2 = \frac{1}{4}$ and $\xi_3 = \frac{1}{3}$

Consequently, we can get

$$\Delta = 1 - \Gamma(q) \sum_{i=1}^m \frac{\alpha_i \eta^{p_i+q-1}}{\Gamma(p_i+q)} - \sum_{i=1}^m \beta_i \xi_i^{q-1} \approx 0.265299.$$

Then, by direct calculations, we can obtain that

$$\Lambda_1 = \sum_{i=1}^m \frac{\alpha_i \eta^{p_i+2(q-1)}}{\Delta \Gamma(p_i+q)} \left(\frac{p_i+q(1-\eta)}{q(p_i+q)} \right) + (q-1) \sum_{i=1}^m \frac{\beta_i (\xi_i^{q-1} - \xi_i^q) \eta^{q-1}}{\Delta} \times \frac{\Gamma(q+1)}{\Gamma(2q+1)} \approx 0.45478$$

$$\Lambda_2 = \left(1 + \frac{\sum_{i=1}^m \beta_i}{\Delta} \right) \frac{\Gamma(q)}{\Gamma(2q)} + \sum_{i=1}^m \frac{\alpha_i \eta^{p_i+q-1}}{\Delta \Gamma(p_i+q)} \left(\frac{p_i+q(1-\eta)}{q(p_i+q)} \right) \approx 2.63219.$$

Choose $r_1 = 1$, $r_2 = 20$, $M_1 = 3$ and $M_2 = 0.35$, $f(t, u)$ satisfies

$$f(t, u) \geq 4 \geq 3 = M_1 r_1, \quad \forall (t, u) \in [0, 1] \times [0, 1]$$

and

$$f(t, u) \leq 6.5 \leq 7 = M_2 r_2 \quad \forall (t, u) \in [0, 1] \times [1, 20]$$

Thus, (H_1) and (H_2) hold. By Theorem 3.1, we have that boundary value problem (4.1) has at least one positive solution u such that $1 < \|u\| < 20$.

4.2. Example. Consider the following boundary value problem:

$$\begin{cases} D^{\frac{3}{2}}u(t) + f(t, u(t)) = 0, & 0 < t < 1, \\ u(0) = 0 \\ u(1) = \frac{1}{8} \left(I^{\frac{1}{2}}u \right) \left(\frac{1}{8} \right) + \frac{1}{3} \left(I^{\frac{3}{2}}u \right) \left(\frac{1}{8} \right) + \frac{1}{4} \left(I^{\frac{5}{2}}u \right) \left(\frac{1}{8} \right) + \frac{1}{3}u \left(\frac{1}{2} \right) + \frac{1}{5}u \left(\frac{1}{8} \right) + \frac{1}{7}u \left(\frac{1}{6} \right), \end{cases} \quad (4.2)$$

where

$$f(t, u) \begin{cases} u \left(\frac{3}{4} - u \right) + \frac{3}{16} (t^2 + 2), & , 0 \leq t \leq 1, 0 \leq u \leq \frac{3}{4}, \\ \frac{1}{4} (t^2 + 2) \cos^2 \left(\frac{2\pi}{9}u \right) + 75 \left(\frac{3}{4} - u \right)^2, & , 0 \leq t \leq 1, \frac{3}{4} \leq u \leq \frac{3}{2}, \\ \frac{1}{16} (t^2 + 677) + \sin^2 \left(u - \frac{3}{2} \right) \pi, & , 0 \leq t \leq 1, \frac{3}{2} \leq u \leq \infty. \end{cases}$$

Set $m = 3, \eta = \frac{1}{8}, q = \frac{3}{2}, \alpha_1 = \frac{1}{8}, \alpha_2 = \frac{1}{3}, \alpha_3 = \frac{1}{4}, p_1 = \frac{1}{2}, p_2 = \frac{3}{2}, p_3 = \frac{5}{2}, \beta_1 = \frac{1}{3}, \beta_2 = \frac{1}{5}, \beta_3 = \frac{1}{7}, \xi_1 = \frac{1}{2}, \xi_2 = \frac{1}{4}$ and $\xi_3 = \frac{1}{6}$.
 Consequently, we can get

$$\Delta = 1 - \Gamma(q) \sum_{i=1}^m \frac{\alpha_i \eta^{p_i+q-1}}{\Gamma(p_i+q)} - \sum_{i=1}^m \beta_i \xi_i^{q-1} \approx 0.689554.$$

Then, by direct calculations, we can obtain that

$$\Lambda_2 = \left(1 + \frac{\sum_{i=1}^m \beta_i}{\Delta} \right) \frac{\Gamma(q)}{\Gamma(2q)} + \sum_{i=1}^m \frac{\alpha_i \eta^{p_i+q-1}}{\Delta \Gamma(p_i+q)} \left(\frac{p_i+q(1-\eta)}{q(p_i+q)} \right) \approx 0,89554,$$

$$\Lambda_3 = \sum_{i=1}^m \frac{\alpha_i \eta^{p_i+2(q-1)} (1-\eta)^q}{\Delta \Gamma(p_i+q) q} + (q-1) \sum_{i=1}^m \frac{\beta_i (\xi_i^{q-1} - \xi_i^q) \eta^{q-1} (1-\eta)^{2q} \Gamma(q+1)}{\Delta \Gamma(2q+1)} \approx 0,030807085.$$

Choose $a = \frac{3}{4}, b = \frac{3}{2}$ and $c = 42.5$, then $f(t, u)$ satisfies

$$f(t, u) \leq \frac{45}{64} < 0.77063113 \approx \Lambda_2^{-1} a, \quad \forall (t, u) \in [0, 1] \times \left[0, \frac{3}{4} \right],$$

$$f(t, u) \geq \frac{43329}{1024} > 39.4002257 \approx \Lambda_3^{-1} b, \quad \forall (t, u) \in \left[\frac{1}{8}, 1 \right] \times \left[\frac{3}{2}, 42.5 \right]$$

and

$$f(t, u) \leq \frac{347}{8} < 43,6690972 \approx \Lambda_2^{-1} c, \quad \forall (t, u) \in [0, 1] \times [0, 42.5].$$

Thus, $(H_3), (H_4)$ and (H_5) hold. By Theorem 3.2, we have that boundary value problem (4.2) has at least one nonnegative solution u_1 and two positive solutions u_2, u_3 such that $\|u_1\| < \frac{3}{4}, \frac{3}{2} < \min_{\frac{1}{8} \leq t \leq 1} u_2(t)$ and $a < \|u_3\|$ with $\min_{\frac{1}{8} \leq t \leq 1} u_3(t) < \frac{3}{2}$.

REFERENCES

- [1] Agarwal, R. P., Alsaedi, A., Alsharif, A., Ahmad, B., On nonlinear fractional-order boundary value problems with nonlocal multi-point conditions involving Liouville-Caputo derivatives, *Differ. Equ. Appl.*, Volume 9, Number 2 (2017), 147–160, doi:10.7153/dea-09-12.
- [2] Bouteraa, N., Benaicha, S., Djourdem, H., Positive solutions for nonlinear fractional differential equation with nonlocal boundary conditions, *Universal Journal of Mathematics and Applications.*, 1 (1) (2018), 39–45.
- [3] Bouteraa, N., Benaicha, S., Existence of solutions for three-point boundary value problem for nonlinear fractional differential equations, *Bulletin of the Transilvania University of Brasov, Series III: Mathematics, Informatics, Physics.*, Vol 10(59), No. 1 2017.
- [4] Cabada, A., Wang, G., Positive solutions of nonlinear fractional differential equations with integral boundary value conditions. *J. Math. Anal. Appl.*, 389 (2012) 403–411.
- [5] Cui, Y., Zou, Y., Existence of solutions for second-order integral boundary value problems. *Nonlinear Anal., Model. Control.*, 21(2016), 828–838.
- [6] Delbosco, D., Fractional calculus and function spaces, *J. Fract. Calc.*, 6 (1996), 45–53.
- [7] Diethelm, K., Freed, D.: On the solutions of nonlinear fractional order differential equations used in the modelling of viscoplasticity. In: Keil, F, Mackens, W, Voss, H, Werthers, J (eds.) *Scientific Computing in Chemical Engineering II - Computational Fluid Dynamics, Reaction Engineering and Molecular Properties*. Springer, Heidelberg., (1999).
- [8] Djourdem, H., S. Benaicha, S., Existence of positive solutions for a nonlinear three-point boundary value problem with integral boundary conditions. *Acta Math. Univ. Comenianae.*, Vol 87, 2 (2018) pp 167–177.
- [9] Glockle, WG., Nonnenmacher, TF., A fractional calculus approach of self-similar protein dynamics. *Biophys. J.*, 68(1995), 46–53.
- [10] Guo, D., V. Lakshmikantham, V., *Nonlinear Problems in Abstract Cones*. Academic Press, New York., (1988).

- [11] Guo, L., Liu, L., Wu, Y., Existence of positive solutions for singular fractional differential equations with infinite-point boundary conditions. *Nonlinear Analysis: Modelling and Control*, Vol. 21, No. 5 2016, 635-650.
- [12] He, J., Some applications of nonlinear fractional differential equations and their approximations. *Bull. Am. Soc. Inf. Sci. Technol.*, **15**(1999), 86-90 .
- [13] Henderson, J., Luca, R., Existence and multiplicity for positive solutions of a system of higher-order multi-point boundary value problems. *Nonlinear Differ. Equ. Appl.*, **20** (2013), 1035-1054
- [14] Henderson, J., Luca, R., Systems of RiemannLiouville fractional equations with multi-point boundary conditions. *Appl. Math. Comput.*, **309** (2017), 303-323.
- [15] Henderson, J., Luca, R., Tudorache, A., On a system of fractionl differential equations with coupled integral boundary conditions. *Fract. Calc. Appl. Anal.*, **18**(2015), 361-386.
- [16] Kilbas, AA., Srivastava, HM., Trujillo, JJ., *Theory and Applications of Fractional Differential Equations*, Elsevier Science B.V., Amsterdam, 2006.
- [17] Leggett, R.W., Williams, L.R., Multiple positive fixed points of nonlinear operators on ordered Banach spaces, *Indiana Univ. Math. J.*, **28** (1979), 673-688.
- [18] Liu, L., Jiang, J., Wu, Y., The unique solution of a class of sum mixed monotone operator equations and its application to fractional boundary value problems. *J. Nonlinear Sci. Appl.*, **9**(5) (2016), 2943-2958.
- [19] Ma, D-X., Positive solutions of multi-point boundary value problem of fractional differential equation, *Arab Journal of Mathematical Sciences.*, vol. 21, no. 2 2015, pp. 225-236.
- [20] Mainardi, F., Fractional calculus: some basic problems in continuum and statistical mechanics. In: Carpinteri, CA, Mainardi, F (eds.) *Fractal and Fractional Calculus in Continuum Mechanics*. Springer., Vienna (1997).
- [21] Metzler, F., Schick, W., Kilian, HG., Nonnenmache, TF., Relaxation in filled polymers: a fractional calculus approach. *J. Chem. Phys.*, **103**(1995), 7180-7186.
- [22] Nyamoradi, N., Existence of solutions for multi-point boundary value problems for fractional differential equations, *Arab J. Math. Sci.*, **18** (2012), 165-175.
- [23] Oldham, K.B., Spanier, J., *The Fractional Calculus*, Academic Press, New York, London, 1974.
- [24] Podlubny, I., *Fractional Differential Equations*, Academic Press, New York, 1999.
- [25] Pu, R.,Zhang, X., Cui, Y., Li, P., Wang, W., Positive solutions for singular semipositone fractional differential equation subject to multipoint boundary conditions, *Journal of Function Spaces.*, vol. 2017, Article ID 5892616, 8 pages, 2017
- [26] Ji, Y.D., Guo, Y.P., Qiu, J.Q., Yang, L.Y., Existence of positive solutions for a boundary value problem of nonlinear fractional differential equations, *Advances in Dierence Equations.*, vol. 2015, article 13, 2015.
- [27] Ross B., (ed.), *The Fractional Calculus and its Applications*, Lecture Notes in Mathematics, Vol. 475, Springer-Verlag, Berlin, 1975.
- [28] Samko, S.G., Kilbas, A.A., Marichev, O.I., *Fractional Integral and Derivatives (Theory and Applications)*, Gordon and Breach, Switzerland, 1993.
- [29] Shah, K., Zeb, S., Khan, R. A., Existence of triple positive solutions for boundary value problem of nonlinear fractional differential equations, *Computational Methods for Differential Equations.*, Vol. 5, No. 2 (2017), pp. 158-169.
- [30] Sun, Y., Zhao, M., Positive solutions for a class of fractional differential equations with integral boundary conditions. *Appl. Math. Lett.*, **34** (2014), 17-21.
- [31] Tariboon, J., Ntouyas, S.K., Sudsutad, W., Positive solutions for fractional differential equations with three-point multi-term fractional integral boundary conditions, *Adv. Difference Equ.*, 2014 (2014), 17 pages.1
- [32] Wang, Y., Liang S., Wang, Q., Multiple positive solutions of fractional-order boundary value problem with integral boundary conditions. *J. Nonlinear Sci. Appl.*, **10** (2017), 6333-6343.
- [33] Webb, J.R.L., Nonlocal conjugate type boundary value problems of higher order. *Nonlinear Anal.*, **71**(2009), 1933-1940.
- [34] Zhou L., Jiang, W., Positive solutions for fractional differential equations with multi-point boundary value problems. *Journal of Applied Mathematics and Physics.*, **2** (2014), 108-114.

HABIB DJOURDEM

LABORATORY OF FUNDAMENTAL AND APPLIED MATHEMATICS OF ORAN (LMFAO), UNIVERSITY OF
ORAN1, AHMED BENBELLA. ALGERIA

E-mail address: `djourdem.habib7@gmail.com`

SLIMANE BENAICHA

LABORATORY OF FUNDAMENTAL AND APPLIED MATHEMATICS OF ORAN (LMFAO), UNIVERSITY OF
ORAN1, AHMED BENBELLA. ALGERIA

E-mail address: `slimanebenaicha@yahoo.fr`