

# Steffensen type methods for approximating solutions of differential equations

Flavius Pătrulescu

**Abstract.** The implicit methods for numerical solving of ODEs lead to nonlinear equations which are usually solved by the Newton method. We study the use of a Steffensen type method instead, and we give conditions under which this method provides bilateral approximations for the solution of these equations; this approach offers a more rigorous control of the errors. Moreover, the method can be applied even in the case when certain functions are not differentiable on the definition domain. The convergence order is the same as for Newton method.

**Mathematics Subject Classification (2010):** 65L04, 65L05, 65H05.

**Keywords:** Initial value problems, stiff equations, Steffensen method, Newton method, convergence order.

## 1. Introduction

The mathematical modeling of many problems in physics, engineering, chemistry, biology, etc. gives rise to ordinary differential equations or systems of ordinary differential equations.

It is known that a high-order initial value problem (IVP) for differential equations or systems of equations can be rewritten as a first-order IVP system (see e.g. [4], [5]) so that the standard IVP can be written in the form:

$$\begin{cases} y' = f(x, y), & x \in I \\ y(a) = y_0, \end{cases} \quad (1.1)$$

where:  $y_0 \in \mathbb{R}^m$ ,  $I \subseteq \mathbb{R}$ ,  $f : I \times \mathbb{R}^m \rightarrow \mathbb{R}^m$  and  $a \in I$ .

A solution is sought on the interval  $[a, b] \subset I$ , where  $a, b$  are finite. In this paper we consider only the scalar case, i.e.,  $m = 1$ .

In practice, the number of cases where an exact solution can be found by analytical means is very limited, so that one uses numerical methods for the approximation of the solution.

Integrating (1.1), for  $m = 1$ , using an implicit linear multistep method with step-size  $h$ , leads to the solving at each step of an equation of the form:

$$y = hA\phi(x, y) + \psi. \quad (1.2)$$

Here  $A$  is a constant determined by the numerical method and  $\psi$  is a known value.

This equation can be solved by the *fixed point iteration*

$$y^{(\nu+1)} = hA\phi(x, y^{(\nu)}) + \psi, \quad y^{(0)} \text{ arbitrary}, \quad \nu = 0, 1, \dots \quad (1.3)$$

which converges to the unique solution of (1.2) provided that:

$$h < \frac{1}{|A|L}, \quad (1.4)$$

where  $L$  is the Lipschitz constant of  $\phi$  with respect to the second variable.

Condition (1.4) becomes too restrictive for stiff problems. Thus, if we use an explicit method to solve a stiff equation, we have to use an excessively small step-size to avoid instability; if we use an implicit method with an absolute stability region large enough to impose no stability restriction, we can choose a step-size as large as we want, but we will not be able to solve the implicit equation (1.2) by the iteration (1.3) unless the step-size is excessively small.

In order to overcome this difficulty one uses the Newton iteration instead of the fixed point iteration. Newton iteration applied to the equation:

$$F(y) = 0, \quad (1.5)$$

where  $F : [c, d] \rightarrow \mathbb{R}$ ,  $c, d \in \mathbb{R}$ ,  $c < d$ , has the form:

$$y^{(\nu+1)} = y^{(\nu)} - F(y^{(\nu)})/F'(y^{(\nu)}), \quad \nu = 0, 1, 2, \dots, \quad y^{(0)} \in [c, d]. \quad (1.6)$$

When applied to the equation (1.2), where  $F(y) = y - hA\phi(x, y) - \psi$ , we get:

$$y^{(\nu+1)} = y^{(\nu)} - (y^{(\nu)} - hA\phi(x, y^{(\nu)}) - \psi)/(1 - hA\phi'_y(x, y^{(\nu)})), \quad (1.7)$$

i.e.,

$$y^{(\nu+1)} = (hA(\phi(x, y^{(\nu)}) - y^{(\nu)}\phi'_y(x, y^{(\nu)})) + \psi)/(1 - hA\phi'_y(x, y^{(\nu)})). \quad (1.8)$$

One step of Newton iteration requests considerably more computing time than one step of fixed point iteration. Each step of the latter costs one function evaluation, whereas each step of the former calls for the updating of the derivative.

In this paper we approximate the solution of equation (1.5) using the Steffensen type method:

$$y^{(\nu+1)} = y^{(\nu)} - \frac{F(y^{(\nu)})}{[y^{(\nu)}, g(y^{(\nu)}); F]}, \quad \nu = 0, 1, \dots \quad (1.9)$$

where  $g : [c, d] \rightarrow [c, d]$  is an auxiliary function such that the equation:

$$y - g(y) = 0 \quad (1.10)$$

is equivalent to (1.5), and  $[u, v; F]$  represents the first order divided difference of  $F$  at the points  $u, v \in [c, d]$ . This method does not require the calculation of the derivative of the function  $F$ .

Let  $y^* \in (c, d)$  be the root of equation (1.5). If the elements of the sequence  $(y^{(\nu)})_{\nu \geq 0}$  belong to the interval  $[c, d]$  then from Newton identity and (1.9) we obtain:

$$y^* - y^{(\nu+1)} = - \frac{[y^*, y^{(\nu)}, g(y^{(\nu)}); F](y^* - y^{(\nu)})(y^* - g(y^{(\nu)}))}{[y^{(\nu)}, g(y^{(\nu)}); F]},$$

where  $[u, v, w; F]$  represents the second order divided difference of  $F$  at the points  $u, v, w \in [c, d]$ . If  $g$  is Lipschitz on  $[c, d]$  with constant  $L$  and if we assume that there exist the real numbers  $M, m > 0$  such that:

$$|[u, v, w; F]| < M \quad \text{and} \quad |[u, v; F]| > m,$$

for all  $u, v, w \in [c, d]$ , then:

$$|y^* - y^{(\nu+1)}| \leq \frac{ML|y^* - y^{(\nu)}|^2}{m},$$

which shows that the  $q$ -convergence order for the method (1.9) is 2, i.e., the same as for the Newton method.

In [7] are given conditions for the convergence of the sequences generated by relation (1.9), and the function  $g$  is defined such that the sequences  $(y^{(\nu)})_{\nu \geq 0}$  and  $(g(y^{(\nu)}))_{\nu \geq 0}$  approximate bilaterally the exact solution  $y^*$ . Thus, we have an a posteriori error control.

For the functions  $F$  and  $g$  we suppose the following hypothesis:

- ( $\alpha$ ) the equations (1.5) and (1.10) are equivalent;
- ( $\beta$ ) the function  $g$  is continuous and decreasing on  $[c, d]$ ;
- ( $\gamma$ ) the equation (1.5) has a unique solution  $y^* \in (c, d)$ .

The following theorem holds (see [7]):

**Theorem 1.1.** *If the functions  $F$  and  $g$  satisfy the conditions ( $\alpha$ ) – ( $\gamma$ ) and moreover the following conditions hold:*

- (i)  $F$  is increasing and convex on  $[c, d]$ ;
- (ii)  $F(y_0) < 0$ ;
- (iii)  $g(y_0) \leq d$ ,

*then the elements of the sequences  $(y^{(\nu)})_{\nu \geq 0}$  and  $(g(y^{(\nu)}))_{\nu \geq 0}$  belong to the interval  $[c, d]$  and the following properties hold:*

- (j) the sequence  $(y^{(\nu)})_{\nu \geq 0}$  is increasing and convergent;
- (jj) the sequence  $(g(y^{(\nu)}))_{\nu \geq 0}$  is decreasing and convergent;
- (jjj)  $y^{(\nu)} \leq y^* \leq g(y^{(\nu)})$ ,  $\nu = 0, 1, \dots$
- (jv)  $\lim_{\nu \rightarrow \infty} y^{(\nu)} = \lim_{\nu \rightarrow \infty} g(y^{(\nu)}) = y^*$ ;
- (vj)  $|y^* - y^{(\nu)}| \leq |g(y^{(\nu)}) - y^{(\nu)}|$ ,  $\nu = 0, 1, \dots$  .

In the above theorem the auxiliary function  $g$  can be taken as:  $g(y) = y - \frac{F(y)}{F'(c)}$ . Similar results have been obtained in [7] if  $F$  verifies the properties:

- F is increasing and concave;  $g$  can be taken as  $g(y) = y - \frac{F(y)}{F'(d)}$ ;
- F is decreasing and concave;  $g$  can be taken as  $g(y) = y - \frac{F(y)}{F'(c)}$ ;
- F is decreasing and convex;  $g$  can be taken as  $g(y) = y - \frac{F(y)}{F'(d)}$ .

The interval  $[a, b]$  is partitioned by the point set  $\{x_n\}$  defined by  $x_n = a + nh$ ,  $n = 0, 1, \dots, N$ ,  $h = (b - a)/N$ , and  $y_n$  denotes an approximation to the exact solution  $y$  of (1.1) at  $x_n$ .

If we use an implicit linear multistep method then  $y_n, n = 1, \dots, N$ , are the solutions of the equation:

$$y = hA\phi(x_n, y) + \psi_n, \tag{1.11}$$

where  $\psi_n = \psi_n(a, h, y_{n-1}, y_{n-2}, \dots, y_0)$ . We call this equation as *approximant equation* and we denote by  $y_n^* \in (c, d), n = 1, \dots, N$ , the exact solution.

For each  $n = 1, \dots, N$  let  $F_n : [c, d] \rightarrow \mathbb{R}$  be defined by

$$F_n(y) = y - hA\phi(x_n, y) - \psi_n. \tag{1.12}$$

Then equation (1.11) can be rewritten in the form  $F_n(y) = 0$ .

To approximate bilaterally the solution  $y_n^*, n = 1, \dots, N$ , we generate the sequence  $(y_n^{(\nu)})_{\nu \geq 0}$ , by:

$$y_n^{(\nu+1)} = y_n^{(\nu)} - \frac{F_n(y_n^{(\nu)})}{[y_n^{(\nu)}, g(y_n^{(\nu)}); F_n]}, \quad \nu = 0, 1, \dots \tag{1.13}$$

or, using (1.12),

$$y_n^{(\nu+1)} = \frac{hA(\phi(x_n, y_n^{(\nu)}) - y_n^{(\nu)}[y_n^{(\nu)}, g(y_n^{(\nu)}); \phi(x_n, \cdot)]) + \psi_n}{1 - hA[y_n^{(\nu)}, g(y_n^{(\nu)}); \phi(x_n, \cdot)]}. \tag{1.14}$$

From Theorem 1.1, if  $F_n$  is increasing and convex, and the initial guess  $y_n^{(0)}$  satisfy  $F_n(y_n^{(0)}) < 0$ , then the sequences  $(y_n^{(\nu)})_{\nu \geq 0}, (g(y_n^{(\nu)}))_{\nu \geq 0}$  converge to  $y_n^*$  and we have the inequalities:

$$y_n^{(\nu)} \leq y_n^* \leq g(y_n^{(\nu)}), \quad \nu = 0, 1, \dots \quad .$$

The rest of the cases can be treated in a similar fashion.

## 2. Application to the trapezoidal rule

We consider the *trapezoidal rule* to integrate the initial value problem (1.1), for  $m = 1$ , and the Steffensen method described above to solve the *approximant equation* (1.11).

The *trapezoidal rule* is a 1-step Adams-Moulton method (an implicit method), and for (1.1) is defined by:

$$y_n = y_{n-1} + \frac{h}{2}(f(x_n, y_n) + f(x_{n-1}, y_{n-1})), n = 1, \dots, N.$$

It is known that the trapezoidal rule is an  $A$ -stable method and has order 2 (see [5]).

For any point  $x_n$ ,  $n = 1, \dots, N$ , we have:

$$y_n - \frac{h}{2}f(x_n, y_n) - \frac{h}{2}f(x_{n-1}, y_{n-1}) - y_{n-1} = 0 \quad (2.1)$$

and in this case  $F_n(y) = y - \frac{h}{2}f(x_n, y) - \frac{h}{2}f(x_{n-1}, y_{n-1}) - y_{n-1}$ . Thus, in (1.11) we have  $A = \frac{1}{2}$ ,  $\phi(x_n, y) = f(x_n, y)$  and  $\psi_n = \frac{h}{2}f(x_{n-1}, y_{n-1}) + y_{n-1}$ ,  $n = 1, \dots, N$ .

For simplicity we consider only the autonomous case, i.e.  $f = f(y)$ , and in this case equation (2.1) becomes:

$$y_n - \frac{h}{2}f(y_n) - \frac{h}{2}f(y_{n-1}) - y_{n-1} = 0 \quad (2.2)$$

and  $F_n(y) = y - \frac{h}{2}f(y) - \psi_n$ ,  $\psi_n = \frac{h}{2}f(y_{n-1}) + y_{n-1}$ ,  $n = 1, \dots, N$ .

Using the fact that

$$[u, v; F_n] = 1 - \frac{h}{2}[u, v; f], \text{ for all } u, v \in [c, d], \quad (2.3)$$

and

$$[u, v, w; F_n] = -\frac{h}{2}[u, v, w; f], \text{ for all } u, v, w \in [c, d], \quad (2.4)$$

$n = 1, \dots, N$ , we obtain that the auxiliary function  $g$  can be taken as (see [10]):

$$g(y) = \frac{\frac{h}{2}(f(y) - y[d - \varepsilon, d; f]) + \psi_n}{1 - \frac{h}{2}[d - \varepsilon, d; f]},$$

or

$$g(y) = \frac{\frac{h}{2}(f(y) - y[c, c + \varepsilon; f]) + \psi_n}{1 - \frac{h}{2}[c, c + \varepsilon; f]},$$

where  $\varepsilon$  is sufficiently small such that the exact solution  $y_n^*$  of the equation  $F_n(y_n) = 0$ ,  $n = 1, \dots, N$ , belongs to the interval  $[c + \varepsilon, d - \varepsilon]$ .

For each  $n = 1, \dots, N$  we denote:

$$\psi_{\max}^n = \max\{y_k + \frac{h}{2}f(y_k) | k = 0, \dots, n-1\},$$

$$\psi_{\min}^n = \min\{y_k + \frac{h}{2}f(y_k) | k = 0, \dots, n-1\}.$$

We are lead to the main results of this work:

**Theorem 2.1.** *If the function  $f$ , the step-size  $h$ , and the initial guesses  $y_n^{(0)}$ ,  $n = 1, \dots, N$ , satisfy the following conditions:*

- (i)  $[u, v, w, f] \leq 0$ , for all  $u, v, w \in [c, d]$ ;
- (ii)  $(m \leq [u, v, f] \leq M \leq 0)$ , for all  $u, v \in [c, d]$  or  $(0 \leq m \leq [u, v, f] \leq M)$ , for all  $u, v \in [c, d]$ , and  $h \leq \frac{2}{M}$ ;
- (iii)  $y_n^{(0)} - \frac{h}{2}f(y_n^{(0)}) < \psi_{\min}^n$ ;
- (iv)  $y_n^{(0)}M - f(y_n^{(0)}) \geq \frac{2}{h}[d(M\frac{h}{2} - 1) + \psi_{\max}^n]$ ,

then the elements of the sequences  $(y_n^{(\nu)})_{\nu \geq 0}$ ,  $(g(y_n^{(\nu)}))_{\nu \geq 0}$ ,  $n = 1, \dots, N$ , belong to the interval  $[c, d]$  and the following properties hold:

- (j)  $(y_n^{(\nu)})_{\nu \geq 0}$  is increasing and convergent;
- (jj)  $(g(y_n^{(\nu)}))_{\nu \geq 0}$  is decreasing and convergent;
- (jjj)  $y_n^{(\nu)} \leq y_n^* \leq g(y_n^{(\nu)})$ ,  $\nu = 0, 1, \dots$ ;
- (jv)  $\lim_{\nu \rightarrow \infty} y_n^{(\nu)} = \lim_{\nu \rightarrow \infty} g(y_n^{(\nu)}) = y_n^*$ ;

$$(v) |y_n^* - y_n^{(\nu)}| \leq |g(y_n^{(\nu)}) - y_n^{(\nu)}|, \quad \nu = 0, 1, \dots$$

*Proof.* From (2.3), (2.4) and (i), (ii) we have  $[u, v; F_n] \geq 0$ ,  $[u, v, w; F_n] \geq 0$ ,  $n = 1, \dots, N$ , for all  $u, v, w \in [c, d]$ , and we deduce that  $F_n$  is increasing and convex.

Also, from (iii) and (iv) we obtain that the initial guesses satisfy the inequalities:  $F(y_n^{(0)}) < 0$  and  $g(y_n^{(0)}) \leq d$ ,  $n = 1, \dots, N$ .

Using Theorem 1.1 we deduce that the properties (j) – (v) hold.  $\square$

The following theorems can be proved in a similar manner:

**Theorem 2.2.** *If the function  $f$ , the step-size  $h$ , and the initial guesses  $y_n^{(0)}$ ,  $n = 1, \dots, N$ , satisfy the following conditions:*

- (i)  $[u, v, w, f] \leq 0$ , for all  $u, v, w \in [c, d]$ ;
- (ii)  $0 \leq m \leq [u, v, f] \leq M$ , for all  $u, v \in [c, d]$ ;
- (iii)  $y_n^{(0)} - \frac{h}{2}f(y_n^{(0)}) < \psi_{\min}^n$ ;
- (iv)  $y_n^{(0)}m - f(y_n^{(0)}) \geq \frac{2}{h}[c(m\frac{h}{2} - 1) + \psi_{\max}^n]$ ;
- (v)  $\frac{2}{m} \leq h$ ,

then the elements of the sequences  $(y_n^{(\nu)})_{\nu \geq 0}$ ,  $(g(y_n^{(\nu)}))_{\nu \geq 0}$ ,  $n = 1, \dots, N$ , belong to the interval  $[c, d]$  and the following properties hold:

- (j)  $(y_n^{(\nu)})_{\nu \geq 0}$  is decreasing and convergent;
- (jj)  $(g(y_n^{(\nu)}))_{\nu \geq 0}$  is increasing and convergent;
- (jjj)  $g(y_n^{(\nu)}) \leq y_n^* \leq y_n^{(\nu)}$ ,  $\nu = 0, 1, \dots$ ;
- (jv)  $\lim_{\nu \rightarrow \infty} y_n^{(\nu)} = \lim_{\nu \rightarrow \infty} g(y_n^{(\nu)}) = y_n^*$ ;
- (v)  $|y_n^* - y_n^{(\nu)}| \leq |g(y_n^{(\nu)}) - y_n^{(\nu)}|$ ,  $\nu = 0, 1, \dots$

**Theorem 2.3.** *If the function  $f$ , the step-size  $h$  and the initial guesses  $y_n^{(0)}$ ,  $n = 1, \dots, N$ , satisfy the following conditions:*

- (i)  $[u, v, w, f] \geq 0$ , for all  $u, v, w \in [c, d]$ ;
- (ii)  $(m \leq [u, v, f] \leq M \leq 0)$ , for all  $u, v \in [c, d]$  or  $(0 \leq m \leq [u, v, f] \leq M)$ , for all  $u, v \in [c, d]$ , and  $h \leq \frac{2}{M}$ ;
- (iii)  $y_n^{(0)} - \frac{h}{2}f(y_n^{(0)}) > \psi_{\max}^n$ ;
- (iv)  $y_n^{(0)}M - f(y_n^{(0)}) \leq \frac{2}{h}[c(M\frac{h}{2} - 1) + \psi_{\min}^n]$ ,

then the elements of the sequences  $(y_n^{(\nu)})_{\nu \geq 0}$ ,  $(g(y_n^{(\nu)}))_{\nu \geq 0}$ ,  $n = 1, \dots, N$ , belong to the interval  $[c, d]$  and the following properties hold:

- (j)  $(y_n^{(\nu)})_{\nu \geq 0}$  is decreasing and convergent;
- (jj)  $(g(y_n^{(\nu)}))_{\nu \geq 0}$  is increasing and convergent;
- (jjj)  $g(y_n^{(\nu)}) \leq y_n^* \leq y_n^{(\nu)}$ ,  $\nu = 0, 1, \dots$ ;
- (jv)  $\lim_{\nu \rightarrow \infty} y_n^{(\nu)} = \lim_{\nu \rightarrow \infty} g(y_n^{(\nu)}) = y_n^*$ ;
- (v)  $|y_n^* - y_n^{(\nu)}| \leq |g(y_n^{(\nu)}) - y_n^{(\nu)}|$ ,  $\nu = 0, 1, \dots$

**Theorem 2.4.** *If the function  $f$ , the step-size  $h$  and the initial guesses  $y_n^{(0)}$ ,  $n = 1, \dots, N$ , satisfy the following conditions:*

- (i)  $[u, v, w, f] \geq 0$ , for all  $u, v, w \in [c, d]$ ;
- (ii)  $0 \leq m \leq [u, v, f] \leq M$ , for all  $u, v \in [c, d]$ ;
- (iii)  $y_n^{(0)} - \frac{h}{2}f(y_n^{(0)}) > \psi_{\max}^n$ ;
- (iv)  $y_n^{(0)}m - f(y_n^{(0)}) \leq \frac{2}{h}[d(m\frac{h}{2} - 1) + \psi_{\min}^n]$ ;
- (v)  $\frac{2}{m} \leq h$ ,

then the elements of the sequences  $(y_n^{(\nu)})_{\nu \geq 0}, (g(y_n^{(\nu)}))_{\nu \geq 0}, n = 1, \dots, N$ , belong to the interval  $[c, d]$  and the following properties hold:

- (j)  $(y_n^{(\nu)})_{\nu \geq 0}$  is increasing and convergent;
- (jj)  $(g(y_n^{(\nu)}))_{\nu \geq 0}$  is decreasing and convergent;
- (jjj)  $y_n^{(\nu)} \leq y_n^* \leq g(y_n^{(\nu)}), \nu = 0, 1, \dots$ ;
- (jv)  $\lim_{\nu \rightarrow \infty} y_n^{(\nu)} = \lim_{\nu \rightarrow \infty} g(y_n^{(\nu)}) = y_n^*$ ;
- (v)  $|y_n^* - y_n^{(\nu)}| \leq |g(y_n^{(\nu)}) - y_n^{(\nu)}|, \nu = 0, 1, \dots$  .

### 3. Numerical example

We consider the autonomous initial value problem:

$$\begin{cases} y'(x) = \cos^2(y(x)), & x \in [0, 1], \\ y(0) = 0. \end{cases} \tag{3.1}$$

The exact solution is  $y : [0, 1] \rightarrow \mathbb{R}, y(x) = \arctan(x)$ , and it is plotted in Figure 1(a) with continuous line.

If we use the *trapezoidal rule* to integrate the above initial value problem we must solve for each mesh point  $x_n = nh, n = 1, \dots, N, h = 1/N, N \in \mathbb{N}$ , the nonlinear equation:

$$y_n = y_{n-1} + \frac{h}{2}(\cos^2 y_n + \cos^2 y_{n-1}), \tag{3.2}$$

where  $x_0 = 0$  and we choose  $y_0 = 0$ .

According to the above sections we can write (3.2) in the form

$$F_n(y) = 0,$$

where  $F_n(y) = y - \frac{h}{2} \cos^2(y) - \psi_n$ , and  $\psi_n = y_{n-1} + \frac{h}{2} \cos^2(y_{n-1}), n = 1, \dots, N$ . It is easy to show that equation (3.2) has a unique solution  $y_n^* \in (0, \frac{\pi}{4}), n = 1, \dots, N$ , and we will use a *Steffensen* type method to obtain a numerical approximation,  $\tilde{y}_n$ , for this solution.

From  $F_n'(y) = 1 + \frac{h}{2} \sin(2y) \geq 0$  and  $F_n''(y) = h \cos(2y) \geq 0, y \in [0, \frac{\pi}{4}], n = 1, \dots, N$ , we deduce that  $F_n$  is increasing and convex. Thus, we can define the decreasing function  $g$  as:

$$g(y) = y - \frac{F_n(y)}{F_n'(0)} = y - F_n(y) = \frac{h}{2} \cos^2(y) + \psi_n, \quad n = 1, \dots, N.$$

Also, from Theorem 2.1, choosing for each  $n = 1, \dots, N$  the initial guesses  $y_n^{(0)}$  such that it verifies the conditions (iii) and (iv) we obtain bilateral approximations of the solution  $y_n^*$  and an a posteriori error control.

The numerical solution, obtained with the method described above, for the step size  $h = 0.05$  is also plotted in Figure 1(a) with circle marker. The values of the errors  $\varepsilon_n = |y(x_n) - \tilde{y}_n|$ ,  $n = 1, \dots, N$ , are presented in the following table. They are also plotted in Figure 1(b). We observe a very good agreement when we compare the numerical with the analytical solution.




FIGURE 1. (a) The exact solution (continuous line) and the numerical solution (circle marker). (b) The values of the errors

TABLE 1. The values of the errors

$x_n$	$\varepsilon_n$	$x_n$	$\varepsilon_n$
0.05	0.00002068828052	0.55	0.00010932962999
0.1	0.00004056160110	0.6	0.00010478854028
0.15	0.00005887122616	0.65	0.00009889278012
0.2	0.00007499149969	0.7	0.00009200032239
0.25	0.00008845999793	0.75	0.00008443386474
0.3	0.00009899709371	0.8	0.00007647332226
0.35	0.00010650491201	0.85	0.00006835314397
0.4	0.00011104895260	0.9	0.00006026315926
0.45	0.00011282772086	0.95	0.00005235176793
0.5	0.00011213634983	1	0.00004473048874



## References

- [1] Agratini, O., Chiorean, I., Coman, Gh., Trîmbițaș, R., *Numerical Analysis and Approximation Theory*, Vol. III, Presa Universitară Clujeană, 2002 (in Romanian).
- [2] Butcher, J.C., *Numerical Methods for Ordinary Differential Equations*, John Wiley & Sons, 2003.
- [3] Căținaș, E., *Methods of Newton and Newton-Krylov Type*, Risoprint, Cluj-Napoca, 2007.
- [4] Crouzeix, M., Mignot, A.L., *Analyse numérique des équations différentielles*, Masson, Paris, 1989.
- [5] Lambert, J.D., *Numerical Methods for Ordinary Differential Systems-The Initial Value Problem*, John Wiley & Sons, 1990.
- [6] Matheij, R., Molenaar, J., *Ordinary Differential Equations in Theory and Practice*, SIAM, Philadelphia, 2002.
- [7] Păvăloiu, I., *On the monotonicity of the sequences of approximations obtained by Steffensen's method*, *Mathematica*, **35(58)**(1993), no. 1, 71-77.
- [8] Păvăloiu, I., *Bilateral approximation for the solution of scalar equations*, *Rev. Anal. Numér. Théor. Approx.*, **23**(1994), no. 1, 95-101.
- [9] Păvăloiu, I., *Approximation of the roots of equations by Aitken-Steffensen-type monotonic sequences*, *Calcolo*, **32**(1995), 69-82.
- [10] Păvăloiu, I., *Aitken-Steffensen type methods for nonsmooth functions(III)*, *Rev. Anal. Numér. Théor. Approx.*, **32**(2003), no. 1, 73-77.

Flavius Pătrulescu  
"Babeș-Bolyai" University  
Faculty of Mathematics and Computer Sciences  
1, Kogălniceanu Street  
400084 Cluj-Napoca  
Romania  
e-mail: fpatrulescu@ictp.acad.ro