

# Coefficient estimates for a subclass of analytic functions by Srivastava-Attiya operator

Mostafa Jafari, Ahmad Motamednezad and Ebrahim Analouei Adegani

**Abstract.** In this paper, we investigate bounds of the coefficients for subclass of analytic and bi-univalent functions. The results presented in this paper would generalize and improve some recent works and other authors.

**Mathematics Subject Classification (2010):** 30C45, 30C50.

**Keywords:** Analytic functions, bi-univalent functions, coefficient estimates, Srivastava-Attiya operator, subordination.

## 1. Introduction

Let  $\mathcal{A}$  be a class of functions of the form

$$f(z) = z + \sum_{n=2}^{\infty} a_n z^n, \quad (1.1)$$

which are analytic in the open unit disk  $\mathbb{U} = \{z \in \mathbb{C} : |z| < 1\}$ . Further, let  $\mathcal{S}$  denote the class of functions  $f \in \mathcal{A}$  which are univalent in  $\mathbb{U}$ .

For  $f(z)$  defined by (1.1) and  $h(z)$  defined by

$$h(z) = z + \sum_{n=2}^{\infty} b_n z^n,$$

the Hadamard product  $(f * h)(z)$  of the functions  $f(z)$  and  $h(z)$  defined by

$$(f * h)(z) = z + \sum_{n=2}^{\infty} a_n b_n z^n,$$

In 2007, Srivastava and Attiya [21] (see also Răducanu and Srivastava [18] and Prajapat and Goyal [17]) for the class  $\mathcal{A}$  introduced and investigated linear operator

$\mathcal{J}_\mu^b : \mathcal{A} \rightarrow \mathcal{A}$  that defined in terms of the Hadamard product by

$$\mathcal{J}_\mu^b f(z) = z + \sum_{k=2}^{\infty} \Theta_k a_k z^k,$$

where

$$\Theta_k = \left| \left( \frac{1+b}{k+b} \right)^\mu \right|,$$

and (throughout this paper unless otherwise mentioned ) the parameters  $\mu, b$  are considered as  $\mu \in \mathbb{C}$  and  $b \in \mathbb{C} \setminus \{0, -1, -2, \dots\}$ , (see for more details [20]).

**Remark 1.1.** (1) For  $\mu = 1$  and  $b = v$  ( $v > -1$ ), we get generalized Libera-Bernardi integral operator [19];  
 (2) For  $\mu = \sigma$  ( $\sigma > 0$ ) and  $b = 1$ , we get Jung-Kim-Srivastava integral operator [12].

For each  $f \in \mathcal{S}$ , the Koebe one-quarter theorem [9] ensures that the image of  $\mathbb{U}$  under  $f$  contains a disk of radius  $\frac{1}{4}$ . Hence every function  $f \in \mathcal{S}$  has an inverse  $f^{-1}$ , which is defined by

$$f^{-1}(f(z)) = z \quad (z \in \mathbb{U}),$$

and

$$f(f^{-1}(w)) = w \quad \left( |w| < r_0(f); r_0(f) \geq \frac{1}{4} \right),$$

where

$$g(w) = f^{-1}(w) = w - a_2 w^2 + (2a_2^2 - a_3) w^3 + \dots \tag{1.2}$$

A function  $f \in \mathcal{A}$  is said to be bi-univalent in  $\mathbb{U}$  if both  $f$  and  $f^{-1}$  are univalent in  $\mathbb{U}$ . Let  $\Sigma$  denote the class of bi-univalent functions in  $\mathbb{U}$  given by (1.1).

Recently many researchers have introduced and investigated several interesting subclasses of the bi-univalent function class  $\Sigma$  and they have found non-sharp estimates on the first two Taylor-Maclaurin coefficients  $|a_2|$  and  $|a_3|$  and other problems, see for example, [3, 2, 4, 5, 6, 7, 8, 10, 11, 13, 14, 15, 22, 23, 24].

For two functions  $f$  and  $g$  that are analytic in  $\mathbb{U}$ , we say that the function  $f$  is *subordinate* to  $g$  and write  $f(z) \prec g(z)$ , if there exists a Schwarz function  $\omega$ , that is analytic in  $\mathbb{U}$  with  $\omega(0) = 0$  and  $|\omega(z)| < 1$  such that  $f(z) = g(\omega(z))$  for all  $z \in \mathbb{U}$ .

In particular, if the function  $g$  is univalent in  $\mathbb{U}$ , then  $f(z) \prec g(z)$  if and only if  $f(0) = g(0)$  and  $f(\mathbb{U}) \subseteq g(\mathbb{U})$ .

In this work, we obtain estimates of coefficients for a subclass of bi-univalent functions considered by Selvaraj et al. [20]. The results presented in this paper would generalize and improve some recent works and other authors.

## 2. The subclass $\mathcal{S}_{\Sigma,t}^{\mu,b}(\gamma, \lambda, \phi)$

Throughout this paper, we assume that  $\phi$  is an analytic function with positive real part in the unit disk  $\mathbb{U}$ , satisfying  $\phi(0) = 1$ ,  $\phi'(0) > 0$  and symmetric with respect to the real axis. Such a function has series expansion of the form

$$\phi(z) = 1 + B_1z + B_2z^2 + B_3z^3 + \dots \quad (B_1 > 0), \tag{2.1}$$

Let that  $u(z)$  and  $v(z)$  are Schwarz function in  $\mathbb{U}$  with

$$u(0) = v(0) = 0, \quad |u(z)| < 1, \quad |v(z)| < 1$$

and suppose that

$$u(z) = \sum_{n=1}^{\infty} p_n z^n \quad \text{and} \quad v(z) = \sum_{n=1}^{\infty} q_n z^n \quad (z \in \mathbb{U}). \tag{2.2}$$

Then [16, p. 172]

$$|p_1| \leq 1, \quad |p_2| \leq 1 - |p_1|^2, \quad |q_1| \leq 1, \quad |q_2| \leq 1 - |q_1|^2. \tag{2.3}$$

By (2.1), we get

$$\phi(u(z)) = 1 + B_1 p_1 z + (B_1 p_2 + B_2 p_1^2) z^2 + \dots \quad (z \in \mathbb{U}) \tag{2.4}$$

and

$$\phi(v(w)) = 1 + B_1 q_1 w + (B_1 q_2 + B_2 q_1^2) w^2 + \dots \quad (w \in \mathbb{U}). \tag{2.5}$$

In 2014, Selvaraj et al. [20] introduced subclass of  $\Sigma$  and obtained estimates on the coefficients  $|a_2|$  and  $|a_3|$  for functions in this subclass as follows:

**Definition 2.1.** [20] A function  $f \in \Sigma$  given by (1.1) is said to be in the class  $\mathcal{S}_{\Sigma,t}^{\mu,b}(\gamma, \lambda, \phi)$  if the following conditions are satisfied:

$$1 + \frac{1}{\gamma} \left( \frac{[(1-t)z]^{1-\lambda} (\mathcal{J}_\mu^b f(z))'}{[\mathcal{J}_\mu^b f(z) - \mathcal{J}_\mu^b f(tz)]^{1-\lambda}} - 1 \right) \prec \phi(z),$$

and

$$1 + \frac{1}{\gamma} \left( \frac{[(1-t)w]^{1-\lambda} (\mathcal{J}_\mu^b g(w))'}{[\mathcal{J}_\mu^b g(w) - \mathcal{J}_\mu^b g(tw)]^{1-\lambda}} - 1 \right) \prec \phi(w),$$

where  $|t| \leq 1$  ( $t \neq 1$ );  $\gamma \in \mathbb{C} \setminus \{0\}$ ;  $\lambda \geq 0$ ;  $z, w \in \mathbb{U}$  and  $g$  is given by (1.2).

**Theorem 2.2.** [20] Let the function  $f(z)$  given by (1.1) be in the class  $\mathcal{S}_{\Sigma,t}^{\mu,b}(\gamma, \lambda, \phi)$ . Then

$$|a_2| \leq \frac{|\gamma| B_1 \sqrt{2B_1}}{\sqrt{|\gamma B_1^2 \Lambda(\lambda, t) \Xi(\lambda, t) - 2(B_2 - B_1)[\Lambda(\lambda, t) + 2]^2 \Theta_2^2 + 2\gamma B_1^2 \Upsilon(\lambda, t) \Theta_3|}} \tag{2.6}$$

and

$$|a_3| \leq \frac{B_1 |\gamma|}{\Upsilon(\lambda, t) \Theta_3} + \left( \frac{B_1 |\tau|}{[\Lambda(\lambda, t) + 2] \Theta_2} \right)^2, \tag{2.7}$$

where

$$\Lambda(\lambda, t) = (\lambda - 1)(1 + t), \quad \Upsilon(\lambda, t) = [(\lambda - 1)(1 + t + t^2) + 3]$$

and

$$\Xi(\lambda, t) = [(\lambda - 2)(1 + t) + 4].$$

### 3. Coefficient estimates

In the section, we get that the following theorem which is an refinement of inequalities (2.6) and (2.7).

**Theorem 3.1.** *Let the function  $f(z)$  given by (1.1) be in the class  $\mathcal{S}_{\Sigma,t}^{\mu,b}(\gamma, \lambda, \phi)$ ,  $|t| \leq 1$  ( $t \neq 1$ ),  $\gamma \in \mathbb{C} \setminus \{0\}$  and  $\lambda \geq 0$ . Then*

$$|a_2| \leq \frac{|\gamma|B_1\sqrt{2B_1}}{\sqrt{2B_1[\Lambda(\lambda, t) + 2]^2\Theta_2^2 + |\gamma B_1^2\Lambda(\lambda, t)\Xi(\lambda, t) - 2B_2[\Lambda(\lambda, t) + 2]^2\Theta_2^2 + 2\gamma B_1^2\Upsilon(\lambda, t)\Theta_3|}}$$

and

$$|a_3| \leq \begin{cases} \frac{|\tau|B_1}{\Upsilon(\lambda, t)\Theta_3} & B_1 \leq \frac{[(\lambda - 1)(1 + t) + 2]^2\Theta_2^2}{|\gamma|\Theta_3[(\lambda - 1)(1 + t + t^2) + 3]} \\ \frac{\Phi(\Theta_1, \Theta_2, \lambda, t)}{\Psi(\Theta_1, \Theta_2, \lambda, t)\Upsilon(\lambda, t)\Theta_3} & B_1 > \frac{[(\lambda - 1)(1 + t) + 2]^2\Theta_2^2}{|\gamma|\Theta_3[(\lambda - 1)(1 + t + t^2) + 3]}. \end{cases}$$

where

$$\Phi(\Theta_1, \Theta_2, \lambda, t) = |\tau|B_1 |\gamma B_1^2\Lambda(\lambda, t)\Xi(\lambda, t) - 2B_2[\Lambda(\lambda, t) + 2]^2\Theta_2^2 + 2\gamma B_1^2\Upsilon(\lambda, t)\Theta_3| + 2|\gamma|^2\Theta_3\Upsilon(\lambda, t)B_1^3,$$

and

$$\Psi(\Theta_1, \Theta_2, \lambda, t) = 2B_1[\Lambda(\lambda, t) + 2]^2\Theta_2^2 + |\gamma B_1^2\Lambda(\lambda, t)\Xi(\lambda, t) - 2B_2[\Lambda(\lambda, t) + 2]^2\Theta_2^2 + 2\gamma B_1^2\Upsilon(\lambda, t)\Theta_3|.$$

*Proof.* Let  $f \in \mathcal{S}_{\Sigma,t}^{\mu,b}(\gamma, \lambda, \phi)$  and  $g = f^{-1}$ . Then there are analytic functions  $u, v : \mathbb{U} \rightarrow \mathbb{U}$ , with  $u(0) = v(0) = 0$ , given by (2.2) such that

$$1 + \frac{1}{\gamma} \left( \frac{[(1 - t)z]^{1-\lambda}(\mathcal{J}_\mu^b f(z))'}{[\mathcal{J}_\mu^b f(z) - \mathcal{J}_\mu^b f(tz)]^{1-\lambda}} - 1 \right) = \phi(u(z)), \tag{3.1}$$

and

$$1 + \frac{1}{\gamma} \left( \frac{[(1 - t)w]^{1-\lambda}(\mathcal{J}_\mu^b g(w))'}{[\mathcal{J}_\mu^b g(w) - \mathcal{J}_\mu^b g(tw)]^{1-\lambda}} - 1 \right) = \phi(v(w)). \tag{3.2}$$

From (2.4), (2.5), (3.1) and (3.2), we obtain

$$[(\lambda - 1)(1 + t) + 2]\Theta_2 a_2 = \gamma B_1 p_1, \tag{3.3}$$

$$[(\lambda - 1)(1 + t + t^2) + 3]\Theta_3 a_3 + \frac{1}{2}(\lambda - 1)(1 + t)[(\lambda - 2)(1 + t) + 4]\Theta_2^2 a_2^2 = \gamma[B_1 p_2 + B_2 p_1^2], \tag{3.4}$$

$$- [(\lambda - 1)(1 + t) + 2]\Theta_2 a_2 = \gamma B_1 q_1, \tag{3.5}$$

and

$$\begin{aligned} & [(\lambda - 1)(1 + t + t^2) + 3]\Theta_3(2a_2^2 - a_3) \\ & + \frac{1}{2}(\lambda - 1)(1 + t)[(\lambda - 2)(1 + t) + 4]\Theta_2^2 a_2^2 = \gamma[B_1 q_2 + B_2 q_1^2]. \end{aligned} \quad (3.6)$$

From (3.3) and (3.5), we get

$$p_1 = -q_1. \quad (3.7)$$

Adding (3.4) and (3.6), and using (3.7), we have

$$\begin{aligned} & ((\lambda - 1)(1 + t)[(\lambda - 2)(1 + t) + 4]\Theta_2^2 + 2\Theta_3[(\lambda - 1)(1 + t + t^2) + 3])a_2^2 \\ & - 2\gamma B_2 p_1^2 = \gamma B_1(p_2 + q_2). \end{aligned} \quad (3.8)$$

From (3.3), we have

$$\begin{aligned} & (\gamma B_1^2\{(\lambda - 1)(1 + t)[(\lambda - 2)(1 + t) + 4]\Theta_2^2 + 2\Theta_3[(\lambda - 1)(1 + t + t^2) + 3]\} \\ & - 2B_2[(\lambda - 1)(1 + t) + 2]^2\Theta_2^2)a_2^2 = \gamma^2 B_1^3(p_2 + q_2). \end{aligned}$$

By (2.3) and (3.3), we get

$$\begin{aligned} & |(\gamma B_1^2\{(\lambda - 1)(1 + t)[(\lambda - 2)(1 + t) + 4]\Theta_2^2 + 2\Theta_3[(\lambda - 1)(1 + t + t^2) + 3]\} \\ & - 2B_2[(\lambda - 1)(1 + t) + 2]^2\Theta_2^2)a_2^2| \leq |\tau|^2 B_1^3(|p_2| + |q_2|) \\ & \leq 2|\gamma|^2 B_1^3(1 - |p_1|^2) \\ & = 2|\gamma|^2 B_1^3 - 2B_1[(\lambda - 1)(1 + t) + 2]^2\Theta_2^2|a_2|^2. \end{aligned}$$

Therefore,

$$\begin{aligned} & |a_2| \leq \quad (3.9) \\ & \leq \frac{|\gamma|B_1\sqrt{2B_1}}{\sqrt{2B_1[\Lambda(\lambda, t) + 2]^2\Theta_2^2 + |\gamma B_1^2\Lambda(\lambda, t)\Xi(\lambda, t) - 2B_2[\Lambda(\lambda, t) + 2]^2\Theta_2^2 + 2\gamma B_1^2\Upsilon(\lambda, t)\Theta_3|}}, \end{aligned}$$

where

$$\Lambda(\lambda, t) = (\lambda - 1)(1 + t), \quad \Upsilon(\lambda, t) = [(\lambda - 1)(1 + t + t^2) + 3]$$

and

$$\Xi(\lambda, t) = [(\lambda - 2)(1 + t) + 4].$$

Next, in order to find the bound on the coefficient  $|a_3|$ , by subtracting (3.6) from (3.4), and using (3.7), we get

$$\begin{aligned} & 2[(\lambda - 1)(1 + t + t^2) + 3]\Theta_3 a_3 = 2\Theta_3[(\lambda - 1)(1 + t + t^2) + 3]a_2^2 \\ & + \tau B_1(p_2 - q_2). \end{aligned} \quad (3.10)$$

Using (2.3) and (3.7), we have

$$\begin{aligned} & 2[(\lambda - 1)(1 + t + t^2) + 3]\Theta_3|a_3| \\ & \leq |\gamma|B_1(|p_2| + |q_2|) + 2\Theta_3[(\lambda - 1)(1 + t + t^2) + 3]|a_2|^2 \\ & \leq 2|\gamma|B_1(1 - |p_1|^2) + 2\Theta_3[(\lambda - 1)(1 + t + t^2) + 3]|a_2|^2. \end{aligned}$$

From (3.3), we get

$$\begin{aligned} & |\gamma|B_1[(\lambda - 1)(1 + t + t^2) + 3]\Theta_3|a_3| \\ & \leq [|\gamma|\Theta_3[(\lambda - 1)(1 + t + t^2) + 3]B_1 - [(\lambda - 1)(1 + t) + 2]^2\Theta_2^2] |a_2|^2 + |\gamma|^2B_1^2. \end{aligned}$$

From (3.9), for  $[|\gamma|\Theta_3[(\lambda - 1)(1 + t + t^2) + 3]B_1 - [(\lambda - 1)(1 + t) + 2]^2\Theta_2^2] > 0$  we have

$$\begin{aligned} & |\gamma|B_1[(\lambda - 1)(1 + t + t^2) + 3]\Theta_3|a_3| \\ & \leq [|\gamma|\Theta_3[(\lambda - 1)(1 + t + t^2) + 3]B_1 - [(\lambda - 1)(1 + t) + 2]^2\Theta_2^2] \\ & \quad \times \frac{2|\gamma|^2B_1^3}{2B_1[\Lambda(\lambda, t) + 2]^2\Theta_2^2 + |\gamma B_1^2\Lambda(\lambda, t)\Xi(\lambda, t) - 2B_2[\Lambda(\lambda, t) + 2]^2\Theta_2^2 + 2\gamma B_1^2\Upsilon(\lambda, t)\Theta_3} \\ & \quad + |\gamma|^2B_1^2. \end{aligned}$$

Therefore,

$$\begin{aligned} |a_3| & \leq [|\gamma|\Theta_3[(\lambda - 1)(1 + t + t^2) + 3]B_1 - [(\lambda - 1)(1 + t) + 2]^2\Theta_2^2] \\ & \quad \times \frac{2|\gamma|B_1^2}{\Psi(\Theta_1, \Theta_2, \lambda, t)[(\lambda - 1)(1 + t + t^2) + 3]\Theta_3} + \frac{|\gamma|B_1}{[(\lambda - 1)(1 + t + t^2) + 3]\Theta_3}, \end{aligned}$$

where

$$\begin{aligned} \Psi(\Theta_1, \Theta_2, \lambda, t) & = 2B_1[\Lambda(\lambda, t) + 2]^2\Theta_2^2 \\ & \quad + |\gamma B_1^2\Lambda(\lambda, t)\Xi(\lambda, t) - 2B_2[\Lambda(\lambda, t) + 2]^2\Theta_2^2 + 2\gamma B_1^2\Upsilon(\lambda, t)\Theta_3|. \end{aligned}$$

Consequently,

$$|a_3| \leq \begin{cases} \frac{|\gamma|B_1}{[(\lambda - 1)(1 + t + t^2) + 3]\Theta_3} & B_1 \leq \frac{[(\lambda - 1)(1 + t) + 2]^2\Theta_2^2}{|\gamma|\Theta_3[(\lambda - 1)(1 + t + t^2) + 3]} \\ \frac{\Phi(\Theta_1, \Theta_2, \lambda, t)}{\Psi(\Theta_1, \Theta_2, \lambda, t)[(\lambda - 1)(1 + t + t^2) + 3]\Theta_3} & B_1 > \frac{[(\lambda - 1)(1 + t) + 2]^2\Theta_2^2}{|\gamma|\Theta_3[(\lambda - 1)(1 + t + t^2) + 3]}, \end{cases}$$

where

$$\begin{aligned} \Phi(\Theta_1, \Theta_2, \lambda, t) & = |\tau|B_1 |\gamma B_1^2\Lambda(\lambda, t)\Xi(\lambda, t) - 2B_2[\Lambda(\lambda, t) + 2]^2\Theta_2^2 + 2\gamma B_1^2\Upsilon(\lambda, t)\Theta_3| \\ & \quad + 2|\gamma|^2\Theta_3[(\lambda - 1)(1 + t + t^2) + 3]B_1^3. \end{aligned}$$

This completes the proof. □

**Remark 3.2.** Theorem 3.1 is an improvement of the estimates obtained by Selvaraj et al. [20] in Theorem 2.2. For the coefficient  $|a_2|$ , it is clear that

$$\begin{aligned} & \frac{|\gamma|B_1\sqrt{2B_1}}{\sqrt{2B_1[\Lambda(\lambda, t) + 2]^2\Theta_2^2 + |\gamma B_1^2\Lambda(\lambda, t)\Xi(\lambda, t) - 2B_2[\Lambda(\lambda, t) + 2]^2\Theta_2^2 + 2\gamma B_1^2\Upsilon(\lambda, t)\Theta_3}}} \\ & \leq \frac{|\gamma|B_1\sqrt{2B_1}}{\sqrt{|\gamma B_1^2\Lambda(\lambda, t)\Xi(\lambda, t) - 2(B_2 - B_1)[\Lambda(\lambda, t) + 2]^2\Theta_2^2 + 2\gamma B_1^2\Upsilon(\lambda, t)\Theta_3}}}. \end{aligned}$$

On the other hand, for the coefficient  $|a_3|$ , we make the following cases:

(i) For  $B_1 \leq \frac{[(\lambda - 1)(1 + t) + 2]^2 \Theta_2^2}{|\gamma| \Theta_3 [(\lambda - 1)(1 + t + t^2) + 3]}$ , it is clear that

$$\frac{|\gamma| B_1}{\Upsilon(\lambda, t) \Theta_3} \leq \frac{B_1 |\gamma|}{\Upsilon(\lambda, t) \Theta_3} + \left( \frac{B_1 |\tau|}{[\Lambda(\lambda, t) + 2] \Theta_2} \right)^2.$$

(ii) For  $B_1 > \frac{[(\lambda - 1)(1 + t) + 2]^2 \Theta_2^2}{|\gamma| \Theta_3 [(\lambda - 1)(1 + t + t^2) + 3]}$ , it is clear that

$$\frac{\Phi(\Theta_1, \Theta_2, \lambda, t)}{\Psi(\Theta_1, \Theta_2, \lambda, t) \Upsilon(\lambda, t) \Theta_3} \leq \frac{B_1 |\gamma|}{\Upsilon(\lambda, t) \Theta_3} + \left( \frac{B_1 |\tau|}{[\Lambda(\lambda, t) + 2] \Theta_2} \right)^2.$$

**Remark 3.3.** If we set  $\lambda = 0$  in Theorem 3.1, then we get an improvement of the estimates obtained by Selvaraj et al. [20, Corollary 2.1].

**Remark 3.4.** If we set  $\lambda = 1$  in Theorem 3.1, then we get an improvement of the estimates obtained by Selvaraj et al. [20, Corollary 2.2].

**Remark 3.5.** If  $\mathcal{J}_\mu^b f(z)$  be the identity map and  $\lambda = 0$  in Theorem 3.1, then we get an improvement of the estimates obtained by Selvaraj et al. [20, Corollary 2.3].

**Remark 3.6.** If  $\mathcal{J}_\mu^b f(z)$  be the identity map and  $\lambda = 1$  in Theorem 3.1, then we get an improvement of the estimates obtained by Selvaraj et al. [20, Corollary 2.4].

**Remark 3.7.** If  $\mathcal{J}_\mu^b f(z)$  be the identity map and  $\gamma = 1$ ,  $t = 0$  in Theorem 3.1, then we get an improvement of the estimates obtained by Deniz [8, Theorem 2.8].

**Remark 3.8.** If  $\mathcal{J}_\mu^b f(z)$  be the identity map and  $\gamma = 1$ ,  $\lambda = 1$  in Theorem 3.1 is an improvement of the estimates obtained by Ali et al. in [3, Theorem 2.1].

**Remark 3.9.** If we take

$$\phi(z) = \frac{1 + Az}{1 + Bz} = 1 + (A - B)z + (B - A)Bz^2 + \cdots \quad (-1 \leq B < A \leq 1, z \in \mathbb{U})$$

and

$$\varphi(z) = \left( \frac{1 + z}{1 - z} \right)^\alpha = 1 + 2\alpha z + 2\alpha^2 z^2 + \cdots \quad (0 < \alpha \leq 1, z \in \mathbb{U}),$$

which gives  $B_1 = A - B$ ,  $B_2 = (B - A)B$  and  $B_1 = 2\alpha$ ,  $B_2 = 2\alpha^2$ , in Theorem 3.1, then we can deduce interesting results analogous, respectively. Also, for suitable choices the parameter  $\mu$  and  $b$  in Theorems 3.1 and some Remarks above we have an improvement of results involving Libera-Bernardi integral operator [19] and Jung-Kim-Srivastava integral operator [12].

**Acknowledgments.** The authors thank from the Najafabad Branch, Islamic Azad University for their financial support.

## References

- [1] Adegani, E.A., Bulut, S., Zireh, A., *Coefficient estimates for a subclass of analytic bi-univalent functions*, Bull. Korean Math. Soc., **55**(2018), 405-413.
- [2] Adegani, E.A., Cho, N.E., Motamednezhad, A., Jafari, M., *Bi-univalent functions associated with Wright hypergeometric functions*, J. Comput. Anal. Appl., **28**(2020), 261-271.
- [3] Ali, R.M., Lee, S.K., Ravichandran, V., Subramaniam, S., *Coefficient estimates for bi-univalent Ma-Minda starlike and convex functions*, Appl. Math. Lett., **25**(2012), 344-351.
- [4] Aouf, M.K., El-Ashwah, R.M., Abd-Eltawab, A.M., *New subclasses of biunivalent functions involving Dziok-Srivastava operator*, ISRN Math. Anal., (2013), Art. ID 387178.
- [5] Brannan, D.A., Taha, T.S., *On some classes of bi-univalent functions*, Stud. Univ. Babeş-Bolyai Math., **31**(1986), 70-77.
- [6] Bulut, S., *Coefficient estimates for a new subclass of analytic and bi-univalent functions defined by Hadamard product*, J. Complex Anal., (2014), Art. ID 302019.
- [7] Çağlar, M., Orhan, H., Yağmur, N., *Coefficient bounds for new subclasses of bi-univalent functions*, Filomat, **27**(2013), 1165-1171.
- [8] Deniz, E., *Certain subclasses of bi-univalent functions satisfying subordinate conditions*, J. Classical Anal., **2**(2013), 49-60.
- [9] Duren, P.L., *Univalent Functions*, Grundlehren der Mathematischen Wissenschaften, Band 259, Springer-Verlag, New York, Berlin, Heidelberg and Tokyo, 1983.
- [10] Frasin, B.A., Aouf, M.K., *New subclasses of bi-univalent functions*, Appl. Math. Lett., **24**(2011), 1569-1573.
- [11] Jafari, M., Bulboacă, T., Zireh, A., Adegani, E.A., *Simple criteria for univalence and coefficient bounds for a certain subclass of analytic functions*, Commun. Fac. Sci. Univ. Ank. Ser. A1 Math. Stat., **69**(2019), no. 1, 394-412.
- [12] Jung, I.B., Kim, Y.C., Srivastava, H.M., *The Hardy space of analytic functions associated with certain one-parameter families of integral operators*, J. Math. Anal. Appl., **176**(1993), 138-147.
- [13] Lewin, M., *On a coefficient problem for bi-univalent functions*, Proc. Amer. Math. Soc., **18**(1967), 63-68.
- [14] Motamednezhad, A., Bulboacă, T., Adegani, E. A., Dibagar, N., *Second Hankel determinant for a subclass of analytic bi-univalent functions defined by subordination*, Turk. J. Math., **42**(2018), 2798-2808.
- [15] Murugusundaramoorthy, G., Bulboacă, T., *Estimate for initial MacLaurin coefficients of certain subclasses of bi-univalent functions of complex order associated with the Hohlov operator*, Ann. Univ. Paedagog. Crac. Stud. Math., **17** (2018), no. 1, 27-36.
- [16] Nehari, Z., *Conformal Mapping*, McGraw-Hill, New York, NY, USA, 1952.
- [17] Prajapat, J.K., Goyal, S.P., *Applications of Srivastava-Attiya operator to the classes of strongly starlike and strongly convex functions*, J. Math. Inequal., **3**(2009), 129-137.
- [18] Răducanu, D., Srivastava, H.M., *A new class of analytic functions defined by means of a convolution operator involving the Hurwitz-Lerch Zeta function*, Integr. Transf. Spec. funct., **18**(2007), 933-943.
- [19] Reddy, G.L., Padmanaban, K.S., *On analytic functions with reference to the Bernardi integral operator*, Bull. Austral. Math. Soc., **25**(1982), 387-396.



- [20] Selvaraj, C., Babu, O.S., Murugusundaramoorthy, G., *Coefficient estimates of bi-Bazilevič functions of Sakaguchi type based on Srivastava-Attiya operator*, FU Math. Inform., **29**(2014), no. 1, 105-117.
- [21] Srivastava, H.M., Attiya, A., *An integral operator associated with the Hurwitz-Lerch Zeta function and differential subordination*, Integr. Transf. Spec. funct., **18**(2007), 207-216.
- [22] Srivastava, H.M., Mishra, A.K., Gochhayat, P., *Certain subclasses of analytic and bi-univalent functions*, Appl. Math. Lett., **23**(2010), 1188-1192.
- [23] Zireh, A., Adegani, E.A., Bidkham, M., *Faber polynomial coefficient estimates for subclass of bi-univalent functions defined by quasisubordinate*, Math. Slovaca, **68**(2018), 369-378.
- [24] Zireh, A., Adegani, E.A., Bulut, S., *Faber polynomial coefficient estimates for a comprehensive subclass of analytic bi-univalent functions defined by subordination*, Bull. Belg. Math. Soc. Simon Stevin, **23**(2016), 487-504.

Mostafa Jafari  
Department of Mathematics,  
Najafabad Branch, Islamic Azad University  
Najafabad, Iran,  
(Corresponding author)  
e-mail: mostafajafari83@gmail.com

Ahmad Motamednezad  
Faculty of Mathematical Sciences,  
Shahrood University of Technology,  
P.O.Box 316-36155, Shahrood, Iran  
e-mail: a.motamedne@gmail.com

Ebrahim Analouei Adegani  
Faculty of Mathematical Sciences,  
Shahrood University of Technology,  
P.O.Box 316-36155, Shahrood, Iran  
e-mail: analoey.ebrahim@gmail.com